



Contribution ID: 233

Type: **Poster**

IFE/P6-05: Computational Study of the Strong Magnetic Field Generation in Non-Spherical Cone-Guided Implosion

Thursday, 11 October 2012 14:00 (4h 45m)

The magnetic field in non-spherical cone-guided implosion is simulated using temporal evolution equations of the magnetic field coupled with simulated result of 2-D radiation hydrodynamic simulation for Fast Ignition as the first attempt. We have found that the magnetic field is generated by $\text{grad}(\text{Te}) \times \text{grad}(\text{Ne})$ term, and it is compressed by the implosion. In the result, it reaches 5 MG at maximum compression, which was not considered before. Also, high Hall parameter region is appeared between cone tip and core plasma. This magnetic field is strong enough to affect the implosion dynamic and hot electron transport, therefore it should be paid attention in these simulations for Fast Ignition.

Country or International Organization of Primary Author

Japan

Primary author: Mr NAGATOMO, Hideo (Japan)

Co-authors: Dr SUNAHARA, Atsushi (Institute for Laser Technology); Prof. SAKAGAMI, HITOSHI (National Institute for Fusion Science); Prof. AZECHI, Hiroshi (Osaka University, Institute of Laser Engineering); Prof. SHIRAGA, Hiroyuki (Osaka University, Institute of Laser Engineering); Prof. MIMA, KUNIOKI (Graduate School for the Creation of New Photonics Industries); Prof. FUJIOKA, Shinsuke (Osaka University, Institute of Laser Engineering); Dr JOHZAKI, Tomoyuki (Institute for Laser Technology)

Presenter: Mr NAGATOMO, Hideo (Japan)

Session Classification: Poster: P6

Track Classification: IFE - Inertial Fusion Experiments and Theory