

Contribution ID: 377

Type: Poster

## EX/P7-09: Validation Studies of Gyrofluid and Gyrokinetic Predictions of Transport and Turbulence Stiffness Using the DIII-D Tokamak

Friday, 12 October 2012 08:30 (4 hours)

A series of carefully designed validation experiments have been conducted on DIII-D to rigorously test gyrofluid and gyrokinetic predictions of transport and turbulence stiffness in both the ion and electron channels. In the first experiment, the ratio of the volume-averaged electron temperature profile to its value at rho=0.84 was found to be essentially invariant with increased heating over a factor of 3 variation in neutral beam injection (NBI) heating in H-mode plasmas with constant pedestal conditions, while a small increase was observed in the ion temperature ratio. The TGLF [1] transport model reproduces the profile measurements and trends with increased NBI heating, providing significant additional support for the fidelity of TGLF in H-modes. Building off these global studies, experiments that quantified local electron stiffness by varying the deposition of electron cyclotron heating (ECH) about a specified reference radius were performed. Applying this technique to L-mode plasmas with no other external heating, a critical inverse temperature gradient scale length a/L\_Te,crit = 2.0±0.3 (1/L\_Te = -nabla T\_e/T\_e, a=0.6 m) has been identified for the first time in DIII-D. Both TGLF and nonlinear GYRO [2] simulations predict similar levels of near-zero turbulence and transport at or below the experimental critical gradient and experimentally relevant levels above it, and a similar transition is observed in linear growth rate calculations. However, both codes also underpredict the power balance calculations of the electron and ion heat fluxes Q\_e and Q\_i, and increases to both a/L\_Te and a/L\_Ti larger than experimental uncertainties are needed to match the power balance results. These results build upon previous validation studies [3] of electron transport at DIII-D, for which new gyrokinetic modeling results will be reported.

[1] G.M. Staebler, et al., Phys. Plasmas 14 (2007) 055909.

[2] J. Candy and R.E. Waltz, Phys. Rev. Lett. 91 (2003) 45001.

[3] J.C. DeBoo et al., Phys. Plasmas 17 (2010) 056105.

Work supported by the US DOE under DE-FG02-07ER54917, DE-FG02-06ER54871, DE-FC02-04ER54698, DE-FC02-99ER54512, DE-FG02-08ER54984, DE-FG02-89ER53296 and DE-FG02-08ER54999.

## **Country or International Organization of Primary Author**

USA

## Primary author: Mr HOLLAND, Christopher (USA)

**Co-authors:** Ms WHITE, Anne E. (Massachusetts Institute of Technology); Dr PETTY, C. Craig (General Atomics); Dr DOYLE, Edward J. (University of California Los Angeles); Dr STAEBLER, Gary M. (General Atomics); Dr MCKEE, George R. (University of Wisconsin-Madison); Dr WANG, Guiding (University of California Los Angeles); Dr DEBOO, James C. (General Atomics); Dr CANDY, Jeffrey (General Atomics); Dr HILLESHEIM, Jon C. (University of California Los Angeles); Dr ZENG, Lei (University of California Los Angeles); Dr SCHMITZ, Lothar (University of California Los Angeles); Dr SNYDER, Philip B. (General Atomics); Dr WALTZ, Ronald E. (General Atomics); Dr SMITH, Sterling

P. (General Atomics); Dr RHODES, Terry L. (University of California Los Angeles); Ms YAN, Zheng (University of Wisconsin-Madison)

**Presenter:** Mr HOLLAND, Christopher (USA)

Session Classification: Poster: P7

Track Classification: EXC - Magnetic Confinement Experiments: Confinement