



Contribution ID: 457

Type: Poster

EX/P2-14: Non-inductive Current Start-up and Plasma Equilibrium with an Inboard Poloidal Field Null by Means of Electron Cyclotron Waves in QUEST

Tuesday, 9 October 2012 14:00 (4h 45m)

ECW Start-up scenarios in QUEST: Non-inductive current start-up using ECW ($f=8.2$ GHz, $P_{\text{rf}} \ll 140$ kW, O-mode, $N_{\parallel} < 0.4$) has been investigated from a view point of multiple ECR interaction, large up-shift and auto resonance condition ($N_{\parallel} \sim 1$) in QUEST. Due to large up-shift of N_{\parallel} at R_{1ce} , ECCD with mildly relativistic electrons moving in the wave momentum direction can be expected. The start-up rate dI_p/dt of 0.3-0.5 MA/sec was achieved within 0.2 s after rf injection. During this phase rapid build up of the energy spectrum of mildly relativistic electrons is observed. This fact strongly supports the ECR current drive scenario based on the relativistic resonance interaction.

Equilibrium with an inboard poloidal field null:

MHD equilibrium state with an inboard poloidal field null is characterized by high β_p . Parameters are $a=0.27$ m, $R_0=0.79$ m, $\Delta/a=0.4$ and $\epsilon \beta_p=1.5$, respectively. Here Δ denotes the Shafranov shift. The R_s locates at ~ 0.5 m. During the flattop phase, I_p reached to 25 kA with a positive dependence of B_z and could be sustained for ~ 10 s. Loop voltage was less than 10 mV, the line density $0.5E18 \text{ m}^{-3}$ ($<$ the cutoff density) and T_e was measured < 100 eV by Thomson scattering. The value of Shafranov λ was kept ~ 5 . Using observed HX temperature of ~ 50 keV and assumed density of the energetic electrons of $\sim 10\%$ of density β_p (=phot/poloidal mag. pressure) due to energetic electrons is consistent with that in MHD equilibrium. Relaxation oscillations of this equilibrium: High β_p equilibrium oscillates at the frequency of ~ 20 Hz under some conditions for 10 –20 kA. The slow rise and sharp drop in I_p are correlated to changes in density, R_s , shift and the HX energy spectrum emitted from the current carrying electrons. The out-of-phase relation in two energy windows (20-100 keV and 200-400 keV) of HXs suggests that electrons are preferably accelerated against the induced return electric field (~ 20 mV/m) during the I_p rise. Thus, confinement of the energetic electrons affects the stability of this equilibrium.

Country or International Organization of Primary Author

Japan

Primary author: Mr ZUSHI, Hideki (Riam Kyushu University)

Co-authors: Dr HASEGAWA, Makoto (RIAM Kyushu University); Mr ISHIGURO, Masaki (AEES Kyushu University); Ms TASHIMA, Saya (AEES Kyushu University)

Presenter: Mr ZUSHI, Hideki (Riam Kyushu University)

Session Classification: Poster: P2

Track Classification: EXW - Magnetic Confinement Experiments: Wave-plasma interactions; current drive; heating; energetic particles