

Contribution ID: 206

Type: Poster

FTP/P7-12: Self-healing of Radiation Damage by Coupled Motion of Grain Boundaries in Tungsten Divertor Plates under Reactor Conditions

Friday, 12 October 2012 08:30 (4 hours)

Radiation damage by fusion neutrons can significantly degrade material properties. In a fusion reactor, longlasting radiation-induced defects such as vacancies, vacancy clusters, and voids introduce additional nuclear safety complication in terms of trap ites for excessive tritium retention. This is especially true in the divertor/first wall of a tokamak like ITER, where extreme thermal stress is also present. Here we use molecular dynamics simulations to elucidate a self-mitigating mechanism in which the large thermal stress can facilitate the recombination of the neutron-collision-cascade-induced vacancies and interstitials through coupled grain boundary (GB) motion in a bcc tungsten under fusion reactor conditions. Specifically, our simulations reveal that for a number of tungsten GBs, absorbing the fast-moving interstitials can help activate coupled GB motion at reduced mechanical stress; the migrating GB then sweeps up the less-mobile vacancies, facilitating vacancy-interstitial recombination inside the GB. We examine the stress-induced mobility characteristics of a number of GBs in W to investigate the likelihood of this cenario.

Country or International Organization of Primary Author

USA

Primary author: Mr BOROVIKOV, Valery (USA)

Co-authors: Dr VOTER, Arthur (Los Alamos National Laboratory); Dr UBERUAGA, Blas (Los Alamos National Laboratory); Dr PEREZ, Danny (Los Alamos National Laboratory); Dr BAI, Xian-Ming (Los Alamos National Laboratory); Dr TANG, Xian-Zhu (Los Alamos National Laboratory)

Presenter: Mr BOROVIKOV, Valery (USA)

Session Classification: Poster: P7

Track Classification: FTP - Fusion Technology and Power Plant Design