



Contribution ID: 572

Type: **Poster**

EX/6-3Rb: ELM Mitigation with SMBI & CJI Fuelling in HL-2A H-mode Plasmas

Friday, October 12, 2012 8:30 AM (4 hours)

For a tokamak fusion reactor, the mitigation of Edge Localized Mode (ELM) is likely a mandatory requirement to avoid excessive erosion of the divertor targets and yet exploit the benefits of high confinement mode operation [1]. It is reported that pellets could trigger ELM when injected in type-I ELMy H-mode and ELM pacing has been accomplished with small multi-pellets injection [2]. Resonant Magnetic Perturbation (RMP) is another effective method to ELM mitigation that full ELM suppression has been achieved on DIII-D and AUG [3, 4]. Recently, ELM amplitude decrease and frequency increase for a finite duration are observed after the Supersonic Molecular Beam Injection (SMBI) fuelling into H-mode plasmas on HL-2A. The same phenomenon is observed even more significantly after the Cluster Jet Injection (CJI) fuelling.

Country or International Organization of Primary Author

China

Primary author: Mr CHEN, Chengyuan (China)

Co-authors: Prof. FENG, Beibin (SWIP); Mr CHENG, Jun (SWIP); Prof. YAO, Lianghua (SWIP); Prof. YANG, Qingwei (SWIP); Mr ZHONG, Wulu (SWIP); Prof. DUAN, Xuru (SWIP); Prof. ZHOU, Yan (SWIP); Dr SHI, zhongbing (SWIP)

Presenter: Mr CHEN, Chengyuan (China)

Session Classification: Poster: P7

Track Classification: EXC - Magnetic Confinement Experiments: Confinement