

Contribution ID: 61

Type: Poster

EX/P6-02: Internal Amplitude, Structure and Identification of CAEs and GAEs in NSTX

Thursday, 11 October 2012 14:00 (4h 45m)

Fast-ions (e.g. fusion alphas and neutral beam ions) will excite a wide range of instabilities in ITER or a Fusion Nuclear Science Facility device. Among the possible instabilities are high frequency Alfvén eigenmodes (AE) excited through Doppler-shifted cyclotron resonance with beam ions [1]. High frequency AEs cause fast-ion transport [2,3,4], correlate with electron thermal transport [5] and are postulated to contribute to ion heating [6]. These high frequency modes have historically been identified variously as compressional (CAE) or global (GAE) Alfvén eigenmodes, but the identification has not proven conclusive. Identification is essential to understanding the extent of their effect, since the two types of modes have very different effects on resonant particle orbits. The effect on plasma performance of high frequency AEs is investigated in NSTX. This is facilitated by a recently upgraded array of 16 fixed-frequency quadrature reflectometers. Detailed measurements of high frequency AE amplitude and eigenmode structure were obtained in a high power (6 MW), beam-heated Hmode plasma (shot 141398) [7] very similar to those discussed in Ref. [5]. These measurements, which extend from the plasma edge to deep in the core, can be used in modeling the effects of the modes on electron thermal transport. The observed modes are identified by comparison of their frequency and measured toroidal mode numbers with local Alfvén dispersion relations [7]. The modes identified as CAEs have higher frequencies and smaller toroidal mode numbers than the GAEs. Also, they are strongly core localized, in contrast with the GAEs, which also peak toward the plasma center but have much broader radial extent.

Work supported by U.S. DOE Contracts DE-FG02-99ER54527 and DE-AC02-09CH11466.

- [1] W.W. Heidbrink, et al., Nucl. Fusion 46, 324 (2006).
- [2] E. D. Fredrickson, et al., Phys. Plasmas 11, 3653 (2004).
- [3] E. D. Fredrickson, et al., Europhys. Conf. Abstracts 35G, P2.119 (2011).
- [4] E. D. Fredrickson, et al., Proc. of 23rd IAEA FEC, Daejon, Korea, 11-16 Oct. 2010, EXW/P7-06.
- [5] D. Stutman, et al., Phys. Rev. Lett. 102, 115002 (2009).
- [6] D. Gates, et al., Phys. Rev. Lett. 87, 205003 (2001).
- [7] N. A. Crocker, et al., Plasma Phys. Controll. Fusion 15, 105001 (2011).

Country or International Organization of Primary Author

United States of America

Primary author: Mr CROCKER, Neal (USA)

Co-authors: Dr LEBLANC, Benoit P. (Princeton Plasma Physics Laboratory); Dr FREDRICKSON, Eric D. (Princeton Plasma Physics Laboratory); Dr YUH, Howard (Nova Photonics); Dr MENARD, Jonathan E. (Princeton Plasma Physics Laboratory); Dr TRITZ, Kevin (The Johns Hopkins University); Dr PODESTÀ, Mario (Princeton Plasma Physics Laboratory); Dr GORELENKOV, Nikolai N. (Princeton Plasma Physics Laboratory); Dr BELL, Ronald E. (Princeton Plasma Physics Laboratory); Dr KUBOTA, Shigeyuki (University of California - Los Angeles); Dr PEEBLES, William A. (University of California - Los Angeles)

Presenter: Mr CROCKER, Neal (USA)

Session Classification: Poster: P6

Track Classification: EXW - Magnetic Confinement Experiments: Wave–plasma interactions; current drive; heating; energetic particles