



Contribution ID: 66

Type: **Poster**

## **EX/P3-01: Poloidal Variation of High-Z Impurity Density due to Hydrogen Minority ICRH on Alcator C-Mod**

*Wednesday, 10 October 2012 08:30 (4 hours)*

In the Alcator C-Mod tokamak, strong, steady-state variations of molybdenum density within a flux surface are routinely observed in plasmas using hydrogen minority ion cyclotron resonant heating. In/out asymmetries, up to a factor of 2, occur with either inboard or outboard accumulation depending on the major radius of the minority resonance layer. Quantitative comparisons between existing parallel high-Z impurity transport theories and experimental results show good agreement when the resonance layer is on the high-field side (HFS) of the tokamak but disagree substantially for low-field side (LFS) heating. Impurity accumulation on the LFS of a flux surface can be explained by the centrifugal force, and is the first observation of intrinsic rotation generating an in/out asymmetries. The accumulation of impurity density on the HFS of a flux surface is shown to be driven by a poloidal potential variation sustained by magnetically trapped non-thermal, cyclotron heated minority ions. Parallel impurity transport theory is extended to account for these fast-ion effects and shown to agree with experimentally measured impurity density asymmetries.

### **Country or International Organization of Primary Author**

USA

**Primary author:** Mr REINKE, Matthew (USA)

**Co-authors:** Dr BADER, Aaron (MIT-PSFC (now at Univ. Wisconsin)); Dr HUBBARD, Amanda (MIT-PSFC); Prof. HUTCHINSON, Ian (MIT-PSFC); Dr HUGHES, Jerry (MIT-PSFC); Dr RICE, John (MIT PSFC); Mr HOWARD, Nathan (MIT-PSFC); Dr DAVID, Pace (MIT-PSFC (now General Atomics)); Dr WUKITCH, Steve (MIT-PSFC); Dr LIN, Yijun (MIT Plasma Science and Fusion Center); Mr PODPALY, Yuri (MIT-PSFC)

**Presenter:** Mr REINKE, Matthew (USA)

**Session Classification:** Poster: P3

**Track Classification:** EXC - Magnetic Confinement Experiments: Confinement