

Contribution ID: 212

Type: Poster

EX/P7-02: Response of Electron-scale Turbulence and Thermal Transport to Continuous ExB Shear Ramping-up in NSTX

Friday, 12 October 2012 08:30 (4 hours)

Microturbulence is considered to be a major candidate in driving anomalous transport in fusion plasmas, and the equilibrium ExB shear generated by externally driven flow can be a powerful tool to control microturbulence in future fusion devices such as FNSF and ITER. Here we present the first observation of the change in electron-scale turbulence wavenumber spectrum (measured by a high-k scattering system) and thermal transport responding to continuous ExB shear ramping-up in an NSTX center-stack limited and NBI-heated L-mode plasma. It is found that while linear stability analysis shows that the maximum ETG mode linear growth rate far exceeds the observed ExB shearing rate in the measurement region of the high-k scattering system, the unstable ITG modes are susceptible to ExB shear stabilization. We observed that as the ExB shearing rate and the maximum ITG mode growth rate continuously increases (from about 0.2 to 0.7) and the maximum power of the measured electron-scale turbulence wavenumber spectra decreases. Meanwhile, both the electron and ion thermal transports are also reduced as long as MHD activities are not important. These observations are found to be consistent with ExB shear stabilization of the ion-scale turbulence.

Country or International Organization of Primary Author

USA

Primary author: Mr REN, Yang (USA)

Co-authors: Dr DOMIER, Calvin (University of California at Davis); Dr MAZZUCATO, Ernesto (Princeton Plasma Physics Laboratory); Dr LEE, K.C. (University of California at Davis); Dr KAYE, Stanley (Princeton Plasma Physics Laboratory); Dr GUTTENFELDER, Walter (Princeton Plasma Physics Laboratory)

Presenter: Mr REN, Yang (USA)

Session Classification: Poster: P7

Track Classification: EXC - Magnetic Confinement Experiments: Confinement