



Contribution ID: 55

Type: **Oral Presentation**

## **TH/4-2: Nonlinear Excitations of Zonal Structures by Toroidal Alfvén Eigenmodes**

*Thursday, October 11, 2012 11:45 AM (20 minutes)*

Zonal flows and, more generally, zonal structures are known to play important self-regulatory roles in the dynamics of microscopic drift-wave type turbulences. Since Toroidal Alfvén Eigenmode (TAE) plays crucial roles in the Alfvén wave instabilities in burning fusion plasmas, it is, thus, important to understand and assess the possible roles of zonal flow/structures on the nonlinear dynamics of TAE. It is shown that zonal flow/structure spontaneous excitation is possible only near the lower accumulation point of the shear Alfvén wave continuous spectrum, including the proper trapped ion responses, causing the zonal structure to be dominated by the zonal current instead of the usual zonal flow. This work shows that including kinetic thermal ion treatment in the nonlinear simulations of Alfvénic modes is an important ingredient for realistic comparisons with experimental measurements, where the existence of zonal currents has been clearly observed.

### **Country or International Organization of Primary Author**

Italy

**Primary author:** Mr ZONCA, Fulvio (Associazione EURATOM-ENEA sulla Fusione)

**Co-author:** Prof. CHEN, Liu (Inst. for Fusion Theory and Simulation, Zhejiang University, Hangzhou 310027, P.R.C. – Dept. of Physics and Astronomy, Univ. of California, Irvine CA 92697-4575, U.S.A.)

**Presenter:** Mr ZONCA, Fulvio (Associazione EURATOM-ENEA sulla Fusione)

**Session Classification:** Waves & Energetic Particles

**Track Classification:** THW - Magnetic Confinement Theory and Modelling: Wave-plasma interactions; current drive; heating; energetic particles