



Contribution ID: 62

Type: Poster

## EX/P6-03: Fast-ion Redistribution and Loss due to Edge Perturbations in the ASDEX Upgrade, DIII-D and KSTAR Tokamaks

*Thursday, 11 October 2012 14:00 (4h 45m)*

The impact of Edge Localized Modes (ELMs) and externally applied resonant and non-resonant Magnetic Perturbations (MPs) on fast-ion confinement / transport has been investigated in the ASDEX Upgrade, DIII-D and KSTAR tokamaks. These studies were enabled by coordinated multi-machine experiments and new diagnostic capabilities that provide detailed experimental results of the interaction between energetic particles and instabilities in particle phase-space. New findings include: bursts of fast-ion losses induced by ELMs dominate the losses in H-mode plasmas as measured by fast-ion loss detectors (FILDs) at different toroidal and poloidal positions; in low-collisionality H-modes, ELM and inter-ELM fluctuations in fast-ion losses appear often strongly connected with main ELM properties and edge flows; filamentary fast-ion losses are observed during ELMs suggesting a strong interaction between fast-ions and the instabilities concomitant to the ELM blobs and filaments; large changes in escaping-ion phase-space are observed within single ELMs; during the mitigation / suppression of type-I ELMs by externally applied MPs, the large fast-ion blobs / filaments observed during ELMs are replaced by a loss of fast-ions with a broad-band frequency and an amplitude of up to 6 times higher than the NBI prompt loss signal without MPs; a clear synergy in the overall fast-ion transport is observed between edge perturbations and internal MHD fluctuations such as Alfvén Eigenmodes (AEs) and sawteeth. Measured fast-ion losses are typically on banana orbits that explore the entire pedestal / Scrape-Off-Layer (SOL). A systematic study of the observed fast-ion losses induced ELMs (mitigated and not-mitigated) and MPs as a function of the MP configuration and magnitude will be presented together with F3D-OFMC and ASCOT simulations of the MP induced fast-ion losses.

### Country or International Organization of Primary Author

Germany

### Collaboration (if applicable, e.g., International Tokamak Physics Activities)

multi-machine contribution: AUG, DIII-D and KSTAR

**Primary author:** Mr GARCIA-MUNOZ, Manuel (Germany)

**Co-authors:** Mr GEIGER, B. (IPP); Dr PACE, D. (General Atomics); Dr WOLFRUM, E. (IPP); Dr CONWAY, G. D. (IPP); Dr POKOL, G. I. (Budapest University of Technology); Dr PARK, H. (POSTECH); Dr CLASSEN, I. G. J. (FOM Institute DIFFER); Mr BOOM, J. (FOM Institute DIFFER); Dr KIM, J. (NFRI); Dr KIM, J. Y. (NFRI & University of Science and Technology - Daejeon); Dr SHINOHARA, K. (JAEA); Dr VAN ZEELAND, M. A. (General Atomics); Dr MARASCHEK, M. (IPP); Mr WILLENSDORFER, M. (Technische Universität Wien); Ms LAZANYI, N. (Budapest University of Technology); Dr LUHMAN, N. (University of California - Davis); Mr ASUNTA, O.

(Aalto University); Dr FISHER, R. K. (General Atomics); Mr AKASLAMPOLO, S. (Aalto University); Dr HAPPEL, T. (IPP); Dr KURKI-SUONIO, T. (Aalto University); Dr RHODES, T. L. (University of California - Los Angeles); Dr LUNT, T. (IPP); Dr IGOCHINE, V. (IPP); Dr SUTTROP, W. (IPP); Prof. HEIDBRINK, W. W. (University of California - Irvine); Dr CHEN, X. (University of California - Irvine)

**Presenter:** Mr GARCIA-MUNOZ, Manuel (Germany)

**Session Classification:** Poster: P6

**Track Classification:** EXW - Magnetic Confinement Experiments: Wave-plasma interactions; current drive; heating; energetic particles