24th IAEA Fusion Energy Conference - IAEA CN-197



Contribution ID: 391 Type: Poster

EX/P8-10: Stabilization of Energetic-Ion-Driven MHD Mode by ECCD in Heliotron J

Friday, 12 October 2012 14:00 (4h 45m)

Second harmonic electron cyclotron current drive (ECCD) has been applied in the Stellarator/Heliotron device, Heliotron J, to control rotational transform profile. Localized EC driven current at central region changes the rotational transform, making a high magnetic shear. An energetic-ion-driven MHD mode of 80 kHz has been completely stabilized when the counter EC current of more than 2 kA is driven. An experiment of scanning the EC driven current shows that there is a threshold in magnetic shear to stabilize the energetic-ion-driven MHD mode.

Country or International Organization of Primary Author

Japan

Primary author: Mr NAGASAKI, Kazunobu (Japan)

Co-authors: Prof. SANO, Fumimichi (Institute of Advanced Energy, Kyoto University); Dr MOTOJIMA, Gen (National Institure for Fusion Science); Dr OKADA, Hiroyuki (Institute of Advanced Energy, Kyoto University); Mr LEE, Hyung Yong (Graduate School of Energy Science, Kyoto University); Dr MASUDA, Kai (Institute of Advanced Energy, Kyoto University); Mr SAKAMOTO, Kinzo (Institute of Advanced Energy, Kyoto University); Dr MUKAI, Kiyofumi (Graduate School of Energy Science, Kyoto University); Dr HANATANI, Kiyoshi (Institute of Advanced Energy, Kyoto University); Mr MIZUNO, Koji (Graduate School of Energy Science, Kyoto University); Mr ZANG, Linge (Graduate School of Energy Science, Kyoto University); Dr MARUSHCHENKO, Nikolai (Max-Planck Institute for Plasma Physics); Dr YAMAMOTO, Satoshi (Institute of Advanced Energy, Kyoto University); Dr KONOSHIMA, Shigeru (Institute of Advanced Energy, Kyoto University); Dr KOBAYASHI, Shinji (Institute of Advanced Energy, Kyoto University); Dr OHSHIMA, Shinsuke (Institute of Advanced Energy, Kyoto University); Dr MINAMI, Takashi (Institute of Advanced Energy, Kyoto University); Mr MINAMI, Takayuki (Graduate School of Energy Science, Kyoto University); Mr KAGAWA, Tasuku (Graduate School of Energy Science, Kyoto University); Dr ESTRADA, Teresa (CIEMAT); Dr MIZUUCHI, Tohru (Institute of Advanced Energy, Kyoto University); Dr YOSHIMURA, Yasuo (National Institute for Fusion Science); Ms NAGAE, Yoko (Graduate School of Energy Science, Kyoto University); Mr NAKAMURA, Yuichi (Graduate School of Energy Science, Kyoto University); Prof. NAKA-MURA, Yuji (Graduate School of Energy Science, Kyoto University); Dr TURKIN, Yuri (Max-Planck Institute for Plasma Physics)

Presenter: Mr NAGASAKI, Kazunobu (Japan)

Session Classification: Poster: P8

Track Classification: EXS - Magnetic Confinement Experiments: Stability