

Contribution ID: 601

Type: Poster

EX/P7-21: Evolution of the Turbulence Radial Wavenumber Spectrum near the L-H Transition in NSTX Ohmic Discharges

Friday, 12 October 2012 08:30 (4 hours)

The measurement of radially extended meso-scale structures such as zonal flows and streamers, as well as the underlying microinstabilities driving them, is critical for under-standing turbulence-driven transport in plasma devices [1]. In particular, the shape and evolution of the radial wavenumber spectrum indicate details of the nonlinear spectral energy transfer [2], the spreading of turbulence [3], as well as the formation of transport barriers [4]. In the National Spherical Torus Experiment (NSTX), the FMCW backscattering diagnostic [5] is used to probe the turbulence radial wavenumber spectrum (k_r = 0-22 cm⁻¹) across the outboard minor radius near the L- to H-mode transition in Ohmic discharges. During the L-mode phase, a broad spectral component (k_r⁻²-10 cm⁻¹) extends over a significant portion of the edge-core from R=120 to 155 cm (rho=0.4-0.95). At the L-H transition, turbulence is quenched across the measurable k_r range at the ETB location, where the radial correlation length drops from ⁻¹.5 to 0.5 cm. The k_r spectrum away from the ETB location is modified on a time scale of tens of microseconds, indicating that nonlocal turbulence dynamics are playing a strong role. Close to the L-H transition, oscillations in the density gradient and edge turbulence quenching become highly correlated. These oscillations are also present in Ohmic discharges without an L-H transition, but are far less frequent. Similar behavior is also seen near the L-H transition in NB-heated discharges.

*Work supported by U.S. DOE Contracts DE-FG02-99-ER54527 and DE-AC02-09CH11466.

[1] A. Fujisawa et al., Plasma Phys. Control. Fusion 53, 124015 (2011).

[2] J. Li and Y. Kishimoto, Phys. Plasmas 17, 072304 (2010).

[3] T.S. Hahm et al., Plasma Phys. Control. Fusion 46, A323 (2004).

[4] W.X. Wang et al., Phys. Plasmas 14, 072306 (2007).

[5] S. Kubota et al., 53rd Meeting of the APS Division of Plasma Physics, B04.00003 (2012).

Country or International Organization of Primary Author

USA

Primary author: Mr KUBOTA, Shigeyuki (USA)

Co-authors: Dr DIALLO, Ahmed (PPPL); Dr LEBLANC, Benoit P. (PPPL); Dr BUSH, Charles E. (ORNL); Dr PARK, Jong-Kyu (Princeton Plasma Physics Laboratory); Dr CROCKER, Neal (University of California Los Angeles); Dr BELL, R.E. (PPPL); Dr MAINGI, Rajesh (Oak Ridge National Laboratory); Dr MAQUEDA, Ricardo J. (Nova Photonics); Dr RAMAN, Roger (University of Washington); Dr KAYE, Stanley (Princeton Plasma Physics Laboratory, Princeton University, Princeton NJ, 08543 USA); Dr ZWEBEN, Stewart J. (PPPL); Dr PEEBLES, William A. (UCLA); Dr REN, Yang (Princeton Plasma Physics Laboratory)

Presenter: Mr KUBOTA, Shigeyuki (USA)

Session Classification: Poster: P7

Track Classification: EXC - Magnetic Confinement Experiments: Confinement