International Symposium on Uranium Raw Material for the Nuclear Fuel Cycle: Exploration, Mining, Production, Supply and Demand, Economics and Environmental Issues (URAM-2018)



Contribution ID: 26

Type: POSTER

GEOCHEMICAL AND MINERALOGICAL CHARACTERIZATION OF THE URANIFEROUS PHOSPHATE ROCKS OF THE NAVAY FORMATION, TÁCHIRA STATE, VENEZUELA

Wednesday 27 June 2018 17:00 (1 hour)

INTRODUCTION

Through the CRP IAEA T11007 a geochemical and mineralogical characterization study of the uraniferous phosphate rocks of the Navay Formation, Táchira state, Venezuela. Phosphates deposit associated with sandstones in the top levels of the Navay Fm. (Upper Cretaceous) in southwest of Táchira state, Venezuela, was discovered in early 1978 by radiometric surveys, conducted by Ministry of Energy and Mines by The National Commission of Nuclear Affair (CONAN). This consists of siliceous shales, calcareous shales, uraniferous phosphatic sandstones and, cherts ("ftanites") [1]. To date there have been studies on the feasibility of exploiting to produce phosphate fertilizers. There have been no studies on uranium mining.

METHODS AND RESULTS

Mineralogical analysis.

Petrographic and mineralogical analyzes by X Ray Diffraction (XRD) it was obtained that the main minerals that make are fluorapatite/chlorapatite (Ca5(PO4)3(F,Cl), collophane or carbonate fluorapatite which is a compositional variant of apatite (Ca5(PO4,CO3)F), uranospathite (Al1-x[]x[(UO2)(PO4)]2(H2O)20+3xF1-3x), quartz (SiO2) (4 –88 %), calcite (CaCO3) (10 –35 %), montmorillonite ((Na,Ca)0,3(Al,Mg)2Si4O10(OH)2) (maximum 6 %), and, microcline (KAlSi3O8) (maximum 1,43 % %) mainly.

The mineralogical studies show the uranium to be present in the following forms:

• Fluorapatite -carbonate fluorapatite (28-75%).

• Uranospathite (2-3%).

Chemical analysis.

Chemical analysis was done by several techniques: portable X Ray Fluorescence (pXRF), Total Reflection X-Ray Fluorescence (TRXRF) and, Inductively Coupled Plasma –Atomic Emission Spectroscopy (ICP AES). According to this the background of the concentration of U is 102 ppm, reaching a maximum value of 160 ppm in a calcareous phosphatic sandstone, what is within the average range of U in marine phosphate rocks (50-300 ppm U) [2].

Sub anomalous values were determined statistically Cd (82 ppm), Cu (261 ppm), Zn (268 ppm), Sr (1832 ppm), Zr (510 ppm) and, anomalous values of Cr (1653 ppm). In addition, maximum values of majority, minority and trace elements were detected: MgO (12.40 %), Al2O3 (9.60 %), K2O (3.07 %), Fe2O3 (1.25 %), MnO (0.05 %), TiO2 (0.35 %), S (1.31 %), Cl (1.07 %), V (82 ppm), Ni (2083 ppm), Pb (86 ppm), Co (199 ppm) and, Rb (91 ppm).

According to the chemical composition of the samples, most correspond to phosphate rocks (median 17.25 % and maximum 24.81 % of P2O5), quartz sandstones (median 25.60 % and maximum 88.70 % of SiO2) and phosphate limestones (median 27.88 % and maximum 70.40 % of CaO) [3].

Gamma Spectrometry.

Gamma spectrometry analysis was also done on several samples of the deposit, in which 226Ra was detected, which may be present in the apatite either replacing the Ca2+ (geochemical affinity) or by the radioactive decay of the 238U series. Other isotopes of this decay series were detected (214Pb and 214Bi).

DISCUSSION AND CONCLUSION

Mineralogy composition.

Fluorapatite has been identified as the main phosphate mineral, which in some cases contains carbonate in its structure (CO32-), which is common in this group of minerals where carbonate ions can replace phosphate ions (PO43-) [4]. The U is hosted in the apatite because the U4+ (ionic radio 0.97 Å) can replace the Ca2+ (ionic radio 0.99 Å) [5]. The presence of uranospathite has been detected by XRD, which is a secondary phosphate of Al and U belonging to the autunite group formed as a result of the weathering of primary phosphates in humid environments [6]. The presence of autunite and wavelite is not ruled out, which have been identified in samples from La Lucha River in Táchira state [1].

The mineralogy is typical of marine phosphatic deposits. Some samples in thin sections were studied by petrography, where they determined that the samples correspond to sandstones phosphatizing, limestones phosphatizing and micritization of peloids.

The fossil Orthokarstenia ewaldi of lower Cretaceous period Maastrichtian age, which indicates a shallow marine environment, was identified by means of petrographic analysis, it is a foraminifera associated with this type of environment [7-8]. The Navay deposit is sandy phosphorites type, formed during the lower Cretacic, probably by currents upwelling, which produced the deposit of these sediments. Constitute a lithological facies deposited in shallow waters while cutting lines and is locally Campanian-Maastrichtian age [9].

The Navay Formation is a lateral equivalent of the La Luna Fm. in western Venezuela, which contains black phosphorites and black shales hydrocarbon-generating [10].

Chemical composition.

The chemical analyzes it was obtained that the U and Ca correlates with the P, indicating that the mineral that host is the apatite. There are strong correlations between P - Y (0.72) which may indicate the presence of yttrium in the mineral phosphates, Ca - Mg (calcite), Ca - Sr (substitution of Sr2+ by Ca2+), Si with Al and K (silicates as microcline and clays such as montmorillonite), Fe -S (possibly forming sulphides), Fe-V (associated in detrital oxides and/or in organic matter), S - V (possibly in organic matter), U with V and Ni (probably associated with organic matter, as well as the strong V - Ni correlation, which may be geochemically associated with porphyrins [11-12].

In several samples yttrium (Y) was detected, which is associated with phosphates, this may be due to ionic substitutions of Y3+ (ionic radio 0,93 Å) in the structure of the apatite by Ca2+ (ionic radio 0,99 Å). It is recommended to perform REE analysis using ICP-MS or another analytical technique, since they could not be detected using the techniques used in this research. The REE can be included in the phosphate minerals, due to their geochemical affinity [2].

By calculating elementary relationships V/Cr - V/V+Ni V/Ni, Navay sediments were deposited under oxic conditions, without replacement, using the Ni/Co ratio, several samples indicate that these sediments were deposited under sub-oxic to anoxic conditions [13-14]. This difference may be due to the fact that there were redox changes during sedimentation, as a result of upwelling currents that brought oxygen-poor water from the bottom to the surface. On the other hand, there may be remobilization, that is, secondary dispersion of several of these elements are redox sensitive and under oxidative conditions by weathering, they have been oxidized and/or dispersed in the deposit.

Gamma spectrometry analysis.

According to the analysis of gamma spectrometry of surface samples and cores, a concentration of average activity of 226Ra of 2100 Bq/kg was detected, in addition to other isotopes of the decay series of 238U were detected. The 226Ra was measured (by the 214Pb and 214Bi measure) and 234Th directly through issuance of 63 keV. The low contrast concentrations activity of the isotope 232Th series and 40K not detected by this technique.

This implies that there are low concentrations of thorium in the deposit, what corresponds to geochemical environments of this type, in which the U is mobilized as U6+ (uranyl ion UO22+), fixed in the apatite as U4+ under reducing conditions, while the Th4+ is immobile in superficial environments, for which its concentration is low in this type of deposits.

By the analysis of gamma spectrometry 226Ra was detected, as well as other isotopes of the decay series of the 238U, however, no isotopes of the 232Th or 40K series were detected.

Conclusions.

The mineralogy of Navay Fm. is typical of marine phosphatic deposits with fluorapatite/chlorapatite, collophane or carbonate fluorapatite which is a compositional variant of apatite (Ca5(PO4,CO3)F), uranospathite, quartz and, calcite mainly.

The presence of autunite and wavelite is not ruled out, which have been identified in samples from La Lucha River in Táchira state, in outcrop Navay Fm.

The mineralogical studies show the uranium to be present in the following forms: apatite (28-75%) and, uranospathite (2-3%).

The fossil Orthokarstenia ewaldi of lower Cretaceous period Maastrichtian age, which indicates a shallow marine environment, was identified by means of petrographic analysis

According to chemical analysis the background of the concentration of U is 102 ppm, reaching a maximum value of 160 ppm in a calcareous phosphatic sandstone, so it can be considered a deposit of uranium as unconventional according to the U grade.

Statistically sub anomalous values were determined Cd (82 ppm), Cu (261 ppm), Zn (268 ppm), Sr (1832 ppm), Zr (510 ppm) and, anomalous values of Cr (1653 ppm), these are considered elements of interest because of their association geochemical with uranium and phosphates.

In several samples yttrium (Y) was detected (Y max. 144 ppm), which is associated with phosphates, it is recommended to perform REE analysis using ICP-MS or another analytical technique, since they could not be detected using the techniques used in this research.

By calculating elementary relationships V/Cr - V/V+Ni V/Ni, Navay sediments were deposited under oxic conditions, but using the Ni/Co ratio, several samples indicate that these sediments were deposited under sub-oxic to anoxic conditions, this difference may be due to the fact that there were redox changes during sedimentation, as a result of upwelling currents that brought oxygen-poor water from the bottom to the surface.

REFERENCES

[1] CÁRDENAS, H., Historia de Caso: El descubrimiento e Investigaciones preliminares de los Fosfatos Uraníferos asociados a las Formaciones Navay-Burgüita, sectores Las Tapas–La Lucha-Fila El Toro, Parroquia San Joaquín de Navay, Municipio Libertador, Estado Táchira. Período 1978-1979, Petroquímica de Venezuela S. A. (PE-QUIVEN) (unpublished report) (2007) 1-27 (in spanish).

[2] CUNEY, M., KYSER, K., Recent and not-so-recent developments in uranium deposits and implications for exploration, Short Course Series, Mineralogical Association of Canada, Quebec City, Quebec 39 (2008) 1-259.
[3] BOGGS, S., Petrology of Sedimentray Rocks, 2nd Ed, Cambridge University Press (2009) 1-612.

[4] KNUDSEN, A., GUNTER, M., Sedimentary Phosphorites –An Example: Phosphoria Formation, Southeastern Idaho, U.S.A., Reviews in Mineralogy and Geochemistry 48 1 (2002) 363.

[5] BRUNETON, P., CUNEY, M., Geology of uranium deposits, Uranium for Nuclear Power, Resources, Mining and Transformation to Fuel 1 (2016) 11.

[6] LOCOCK, A., et al., The structure and composition of uranospathite, $Al1-x_x[(UO2)(PO4)]2(H2O)20+3xF1-3x$, 0 < x < 0.33, a non-centrosymmetric fluorine-bearing mineral of the autunite group, and of a related synthetic lower hydrate, $Al0.67_0.33[(UO2)(PO4)]2(H2O)15.5$, The Canadian Mineralogist 43 (2005) 989.

[7] CRUZ, L., et al., Caracterización físico química, taxonomía y ecología de orthokarstenia ewaldi (foraminiferida: siphogenerinoididae) de la Formación Los Pinos (cretácico: maastrichtiano) de Samacá (Boyacá, Colombia), Boletín de Geología 33 2 (2011) 95 (in spanish).

[8] PARRA, M., et al., Late Cretaceous Anoxia and Lateral

Microfacies Changes in the Tres Esquinas Member, La Luna Formation, Western Venezuela, Palaios 18 4-5 (2003) 321.

[9] KISER, G., Review of the Cretaceous stratigraphy of the Barinas mountain front, Boletin A.V. G. M. P. 11 (1961) 335.

[10] APPLETON, J, D., NOTHOLT, A, J, G., Local phosphate resources for sustainable development in central and south America, Economic Minerals and Geochemical Baseline Program, Report CR/02/122/N, British Geological Survey, Natural Environment Research Council, Keyworth, Nottingham (2002) 1-102.

[11] GAO, Y, Y., et al., Vanadium: Global (bio) geochemistry, Chemical Geology 417 (2014) 68.

[12] HUANG, J, H., et al., Distribution of Nickel and Vanadium in Venezuelan Crude Oil, Petroleum Science and Technology 31 (2013) 509.

[13] SÁEZ, R., et al., Black shales and massive sulphide deposits: causal or casual relationships? Insights from Rammelsberg, Tharsis, and Draa Sfar, Miner Deposita 46 (2011) 585.

[14] JONES, B., MANNING, D., Comparison of geochemical indices used for the interpretation of paleoredox conditions in ancient mudstones, Chemical Geology 111 (1994) 111.

Country or International Organization

Ecuador

Author: Mr MANRIQUE, John (Universidad Técnica Particular de Loja)

Co-authors: Mr LINARES, Eduard (Universidad Central de Venezuela); Dr BARROS, Haydn (Universidad Simón Bolívar); Mr MONSALVE, Jesús (Universidad Central de Venezuela)

Presenter: Mr MANRIQUE, John (Universidad Técnica Particular de Loja)

Session Classification: Poster Session

Track Classification: Track 3. Applied geology and geometallurgy of uranium and associated metals