

# Recent advances in ICRF heating of mixture plasmas: survey of JET and AUG experiments and extrapolation to JET-DT and ITER

Friday, October 26, 2018 3:20 PM (20 minutes)

This contribution summarizes recent theoretical and experimental developments of a novel ‘three-ion species’ heating scheme that have opened new promising avenues for the application of ICRF in fusion plasmas. Following successful proof-of-principle demonstration on the Alcator C-Mod and JET tokamaks [1], this scenario has also been recently established on AUG. A small amount of  $^3\text{He}$  ions ( $\sim 1\%$  and below) was injected into H-D plasmas to absorb RF power and heat the plasma. In JET experiments, effective plasma heating was observed both at extremely low  $^3\text{He}$  concentrations of  $\sim 0.1\text{-}0.2\%$  (maximized fast-ion content) and at moderate concentrations of  $\sim 1\text{-}1.5\%$ . We further enhanced the efficiency for fast-ion generation and plasma heating by changing the configuration of ICRH antennas from dipole to  $+\pi/2$  phasing. Heating AUG plasmas with this ICRF scenario requires  $^3\text{He}$  ions to be less energetic than in JET. The combination of moderate  $^3\text{He}$  concentrations of  $\sim 1\%$  and off-axis  $^3\text{He}$  resonance was successfully applied to reduce fast-ion energies and thus improve confinement of RF-heated ions in AUG. ICRH modeling with the state-of-the-art codes SCENIC [2] and TORIC-SSFPQL has been used extensively to validate JET and AUG experimental observations.

In a next-step, we also successfully demonstrated effective heating of JET H-D mixtures using the fast injected D-NBI ions as resonant ‘third’ species [3]. The scenario was tuned such that D-NBI ions with injection energy of 100keV absorbed most of launched RF power and were accelerated with ICRF up to  $\sim 2\text{MeV}$ . The observed ten-fold increase in the neutron rate and its temporal evolution were successfully reproduced with the time-dependent TRANSP modeling. The established technique of accelerating NBI ions in mixture plasmas to higher energies can be applied to generate alpha particles in D- $^3\text{He}$  plasmas and to maximize D-T fusion reactivity.

Finally, we conclude with a discussion of the application of these novel ICRF scenarios for future JET-DT and ITER operations [4].

[1] Ye.O. Kazakov, J. Ongena, J.C. Wright, S.J. Wukitch et al, *Nature Physics* **13**, 973-978 (2017)

[2] J.M. Faustin et al, *Plasma Phys. Control. Fusion* **59**, 084001 (2017)

[3] J. Ongena, Ye.O. Kazakov et al, *EPJ Web Conf.* **157**, 02006 (2017)

[4] M. Schneider, J.-F. Artaud, P. Bonoli, Y. Kazakov et al, *EPJ Web Conf.* **157**, 03046 (2017)

## Country or International Organization

Belgium

## Paper Number

EX/8-1

**Primary author:** Dr KAZAKOV, Yevgen (Laboratory for Plasma Physics, LPP-ERM/KMS, Brussels, Belgium)

**Co-authors:** JET CONTRIBUTORS, - (-); THE ASDEX UPGRADE TEAM, - (-); THE EUROFUSION MST1 TEAM, - (-); Dr CZARNECKA, Agata (IPPLM, Warsaw, Poland); Dr CARDINALI, Alessandro (ENEA, Italy); Dr KAPPATOU, Athina (Max-Planck-Institute for Plasma Physics, Garching, Germany); Dr GIROUD, Carine (CCFE); Dr HELLESEN, Carl (Uppsala University, Sweden); Dr CASTALDO, Carmine (ENEA); Mr GALLART, Daniel (BSC, Barcelona, Spain); Dr VALCARCEL, Daniel (UKAEA); Dr VAN EESTER, Dirk (LPP-ERM/KMS); Dr GREKOV, Dmytro (KIPT, Kharkiv, Ukraine); Dr LERCHE, Ernesto Augusto (LPP-ERM/KMS); Dr NABAIS, Fernando (IST, Lisbon, Portugal); Mr DURODIE, Frederic (LPP-ERM/KMS); Mr PATTEN, Hamish (EPFL, Lausanne, Switzerland); Dr

WEISEN, Henri (EPFL, Lausanne, Switzerland); Dr MONAKHOV, Igor (UKAEA); Dr ERIKSSON, Jacob (Uppsala University, Sweden); Dr BIELECKI, Jakub (Institute of Nuclear Physics, Krakow, Poland); Mr GONZALEZ-MARTIN, Javier (University of Seville, Spain); Prof. NOTERDAEME, Jean-Marie (Max Planck Institute for Plasma Physics); Mr GALDON-QUIROGA, Joaquin (University of Seville); Dr WRIGHT, John (MIT - PSFC); Dr FAUSTIN, Jonathan (Max-Planck-Institute for Plasma Physics, Greifswald, Germany); Dr GRAVES, Jonathan (EPFL, Lausanne, Switzerland); Dr ONGENA, Jozef (Plasma Physics Lab, ERM-KMS, Brussels); Dr CROMBE, Kristel (Ghent University, Belgium); Dr GIACOMELLI, Luca (Istituto di Fisica del Plasma, CNR, Milan, Italy); Dr GARCIA-MUNOZ, Manuel (Max-Planck Institute for Plasma Physics); Dr GONICHE, Marc (CEA, IRFM); Dr NAVE, Maria Filomena (Instituto de Plasmas e Fusão Nuclear, Instituto Superior Técnico); Dr WEILAND, Markus (Max-Planck-Institut für Plasmaphysik); Dr NOCENTE, Massimo (Dipartimento di Fisica, Università di Milano-Bicocca); Dr MANTSINEN, Mervi (BSC, Barcelona Spain; ICREA, Barcelona, Spain); Dr FITZGERALD, Michael (UKAEA); Prof. PORKO-LAB, Miklos (MIT); Dr SCHNEIDER, Mireille (ITER Organization, France); Dr LENNHOLM, Morten (European Commission); Ms KRAWCZYK, Natalia (IPPLM, Warsaw, Poland); Dr SCHNEIDER, Philip A. (Max-Planck-Institut für Plasmaphysik); Dr JACQUET, Philippe (CCFE); Mr DUMORTIER, Pierre (LPP-ERM/KMS); Dr DUMONT, Remi (CEA, France); Dr FELTON, Robert (UKAEA); Dr BILATO, Roberto (Max-Planck-Institute for Plasma Physics, Garching, Germany); Dr OCHOUKOV, Roman (Max-Planck-Institute for Plasma Physics, Garching, Germany); Dr SHARAPOV, Sergei (CCFE); Dr BOZHENKOV, Sergey (Max-Planck Institute of Plasma Physics); Dr MENMUIR, Sheena (UKAEA); Dr WUKITCH, Stephen (MIT PSFC); Dr CRACIUNESCU, Teddy (National Institute for Laser, Plasma and Radiation Physics, Bucharest, Romania); Dr LOARER, Thierry (CEA, IRFM, France); Dr JOHNSON, Thomas (KTH, Stockholm, Sweden); Dr KIPTILY, Vasily (UKAEA); Dr BOBKOV, Volodymyr (Max-Planck-Institute for Plasma Physics, Garching, Germany); Dr BARANOV, Yuriy (UKAEA)

**Presenter:** Dr KAZAKOV, Yevgen (Laboratory for Plasma Physics, LPP-ERM/KMS, Brussels, Belgium)

**Session Classification:** EX/8, PPC/2 - TH/6 Heating, Current Drive & Steady State

**Track Classification:** EXW - Magnetic Confinement Experiments: Wave-plasma interactions; current drive; heating; energetic particles