

Overview of Physics Studies on ASDEX Upgrade

Monday, October 22, 2018 2:00 PM (25 minutes)

The ASDEX Upgrade (AUG) programme, jointly run with the EUROfusion MST1 task force, continues to enhance significantly the physics base of ITER and DEMO. Here, the full tungsten wall is a key asset for extrapolating to future devices. The high overall heating power and flexible heating mix and comprehensive diagnostic set allows studies ranging from mimicking the scrape-off-layer (SOL) and divertor conditions of ITER and DEMO at high density to fully non-inductive operation ($q_95=5.5$, $\beta_N \leq 2.8$) at low density. Higher ECRH heating power ≤ 8 MW, new diagnostics and improved analysis techniques have enhanced the capabilities of AUG. Stable high-density H-modes with $P_{sep}/R \leq 11$ MW/m with fully detached strike-points have been demonstrated. The ballooning instability close to the separatrix has been identified as a potential cause leading to the H-mode density limit. Density limit disruptions have been successfully avoided using a path-oriented approach to disruption handling and progress has been made in understanding the dissipation and avoidance of runaway electron beams. ELM suppression with resonant magnetic perturbations (RMP) is now routinely achieved reaching $HH_{98}(y,2) \leq 1.1$ giving new insight into the field penetration physics, in particular with respect to plasma flows. Modelling agrees well with plasma response measurements and a helically localised ballooning structure observed prior to the ELM is evidence for the changed edge stability due to the RMP. Fast measurements of T_i and E_r show that the dominantly neoclassical character of E_r holds through the ELM recovery. Good agreement of 3D nonlinear MHD modelling with measured ELM crash dynamics is achieved. As type-I ELMs (even mitigated) are likely not a viable operational regime in DEMO studies of no ELM regimes have been extended. Stable I-modes up to $n/n_{GW} \leq 0.7$ have been characterised using beta feedback. Despite the sub-Alfvénic beam energy nonlinear energetic particle modes have been observed allowing modelling comparisons under burning plasma conditions. First measurements of the eddy tilt angle of n_e fluctuations using correlation Doppler reflectometry as well as the radial correlation and cross-phase angles of T_e fluctuations have been achieved, showing good agreement with Gyrokinetic simulations. Dedicated matches of H, D and He discharges (core/edge) highlight important isotope physics.

Country or International Organization

United Kingdom

Paper Number

OV/2-1

Primary author: Dr MEYER, Hendrik (UK Atomic Energy Authority)

Presenter: Dr MEYER, Hendrik (UK Atomic Energy Authority)

Session Classification: OV/2 Overview Magnetic Fusion

Track Classification: OV - Overviews