

## Scenario development for DT operation at JET

Wednesday 24 October 2018 15:40 (20 minutes)

The JET exploitation plan foresees D-T operations in 2019-20. With respect to the first D-T campaign in 1998, when JET was equipped with a C wall, the experiments will be conducted in presence of a Be-W ITER-like wall and will benefit from an extended and improved set of diagnostics and higher available additional power. Among the challenges presented by operations with the new wall there are a general deterioration of the pedestal confinement, the risk of heavy impurity accumulation in the core, and the requirement to protect the W divertor from excessive heat loads. Therefore, an intense activity of scenario development has been undertaken at JET during the last three years to overcome these difficulties and to achieve a stationary scenario of the duration of 5 seconds featuring  $H_{98} > 0.9$ ,  $W_{th} \approx 10-12$  MJ towards the lowest values of  $\rho$  and  $\nu$  achievable on JET.

Two complementary scenarios are being developed to approach the problem of developing a scenario suitable for high-performance D-T operation. The baseline scenario ( $\beta_N \sim 1.8$  and  $H_{98} \sim 1.0$ ) concentrates mainly on pushing the operation towards the high current and field limits with a relaxed current profile, whereas the hybrid scenario ( $\beta_N \sim 2-3$  and  $H_{98} > 1.0$ ) exploits the advantages of operating at high normalised beta with a shaped current profile above unity. Encouraging results were achieved for the baseline scenario at 3MA/2.8T and for the hybrid scenario at reduced plasma current (2.2-2.5MA/2.8-2.9T). High-performance plasmas with  $H_{98} \sim 0.9$  producing  $\sim 3 \cdot 10^{16}$  neutrons/s were obtained for  $> 5$  energy confinement times ( $\sim 1.5$ s).

A third scenario, has also been developed for alpha particle studies. This scenario aims at maintaining high plasma performance for 1-2s to generate a significant population of  $\alpha$ -particle for the  $\alpha$ -particle studies and deliberately omits ICRH heating to avoid creating RF driven fast particles, which could mask the effect of the fusion-generated  $\alpha$ -particles. In these pulses ICRH induced TAEs were observed after the NBI switch-off compatibly with the beam fast ion slowing-down time.

The results of all scenarios have been the object of an extensive activity of code validation and modelling and extrapolated to the target D-T scenarios.

### Country or International Organization

United Kingdom

### Paper Number

EX/3-6

**Author:** Dr GARZOTTI, Luca (United Kingdom Atomic Energy Agency - Culham Centre for Fusion Energy)

**Co-authors:** Dr SIPS, Adrianus (EFDA-JET); Dr CZARNECKA, Agata (IFPILM); Dr GIROUD, Carine (CCFE); Dr LOWRY, Christopher (European Commission); Dr CHALLIS, Clive (Culham Centre for Fusion Energy); Dr VALCARCEL, Daniel (UKAEA); Dr FRIGIONE, Domenico (ENEA); Dr JOFFRIN, Emmanuel (CEA); Dr LERCHE, Ernesto Augusto (LPP-ERM/KMS); Dr STEFANIKOVA, Estera (KTH); Dr RIMINI, Fernanda (UKAEA); Dr CASSON, Francis (UKAEA); Dr KIM, Hyun-Tae (EUROfusion Consortium JET); Dr FERREIRA NUNES, Isabel Maria (IPFN/IST); Dr GARCIA, Jeronimo (CEA IRFM); Dr MAILLOUX, Joelle (UKAEA); Dr GRAVES, Jonathan (EPFL); Dr FRASSINETTI, Lorenzo (KTH, Royal Institute of Technology); Dr MENESES, Luis (IST); Dr TSALAS, Maximos (DIFFER); Dr MANTSINEN, Mervi (BSC); Dr ROMANELLI, Michele (CCFE); Mr LENNHOLM, Morten (European Commission); Dr KRAWCZYK, Natalia (IFPILM); Dr LOMAS, Peter (UKAEA); Dr DUMONT, Remi (CEA); Dr FELTON, Robert (UKAEA); Dr SILBURN, Scott (UKAEA); Dr SHARAPOV, Sergei (CCFE)

**Presenter:** Dr GARZOTTI, Luca (United Kingdom Atomic Energy Agency - Culham Centre for Fusion Energy)

**Session Classification:** EX/3 Plasma Performance & Control

**Track Classification:** EXC - Magnetic Confinement Experiments: Confinement