

Fully Non-inductive 2nd Harmonic Electron Cyclotron Current Ramp-up with Focused Polarized Beams in the QUEST Spherical Tokamak

Wednesday, 24 October 2018 08:30 (4 hours)

A transmission line and a launcher system have been newly developed to conduct the second (2nd) harmonic electron cyclotron (EC) plasma ramp-up with an eXtra-ordinary mode wave in the QUEST spherical tokamak. The incident elliptical polarizations were controlled with two corrugated (quarter/one-eighth wavelength) polarizers. The launcher system with two quasi-optical mirrors produced a sharply focused incident beam with a waist size of 0.05 m at the 2nd electron cyclotron resonance layer. The obtained electron density was one order of magnitude higher, compared to the previous experiments with no polarized focusing-beam. As a new record of non-inductive plasma ramp-up with EC-waves, a highest plasma current of 86 kA was achieved with a focused 230 kW 28GHz-beam. The record plasma current ramp-up efficiency on the incident power in the 2nd harmonic EC scenario was also achieved.

Country or International Organization

Japan

Paper Number

EX/P3-21

Primary author: Dr IDEI, Hiroshi (Research Institute for Applied Mechanics, Kyushu University)

Co-authors: Ms HIGASHIJIMA, Aki (Research Institute for Applied Mechanics, Kyushu University); Dr EJIRI, Akira (Graduate School of Frontier Sciences, The University of Tokyo); Prof. FUKUYAMA, Atsushi (Kyoto University); Dr TAYLOR, Gary (Princeton Plasma Physics Laboratory); Prof. HANADA, Kazuaki (Advanced Fusion Research Center, Research Institute for Applied Mechanics, Kyushu University); Dr NAKAMURA, Kazuo (Research Institute for Applied Mechanics, Kyushu University); Mr NAGATA, Kazuya (Research Institute for Applied Mechanics, Kyushu University); Dr KURODA, Kengoh (Research Institute for Applied Mechanics, Kyushu University); Dr MISHRA, Kishore (Institute for Plasma Research); Dr HASEGAWA, Makoto (Research Institute for Applied Mechanics, Kyushu University); Mr FUKUYAMA, Masaharu (Interdisciplinary Graduate School of Engineering Sciences, Kyushu University); Dr ONO, Masayuki (PPPL/Princeton University); Ms YUNOKI, Miyu (Interdisciplinary Graduate School of Engineering Sciences, Kyushu University); Mr MATSUMOTO, Naoki (University of Tokyo); Dr BERTELLI, Nicola (Princeton Plasma Physics Laboratory); Dr WATANABE, Osamu (Research Institute for Applied Mechanics, Kyushu University); Mr YONEDA, Ryota (Interdisciplinary Graduate School of Engineering Sciences, Kyushu University); Dr MURAKAMI, Sadayoshi (Department Nuclear Engineering, Kyoto University); Mr KOBAYASHI, Sakuji (National Institute for Fusion Science); Prof. KUBO, Shin (National Institute for Fusion Science); Mr KOJIMA, Shinichiro (Interdisciplinary Graduate School of Engineering Sciences, Kyushu University); Dr ONCHI, Takumi (Research Institute for Applied Mechanics, Kyushu University); Dr TSUJIMURA, Toru (National Institute for Fusion Science); Prof. IMAI, Tsuyoshi (Plasma Research Center, University of Tsukuba); Dr KARIYA, Tsuyoshi (Plasma Research Center, University of Tsukuba); Dr NAGASHIMA, Yoshihiko (Research Institute for Applied Mechanics, Kyushu University); Prof. TAKASE, Yuichi (University of Tokyo)

Presenter: Dr IDEI, Hiroshi (Research Institute for Applied Mechanics, Kyushu University)

Session Classification: P3 Posters