

# Effect of poloidal density asymmetries on shear flows and radial electric field at the plasma edge

*Friday, October 26, 2018 2:00 PM (20 minutes)*

In the simplest, magnetohydrodynamic (MHD) description of the plasma equilibrium, rapid transport along the field lines leads to a state where the plasma density and temperature are constant on flux surfaces, exhibiting symmetry in both the poloidal and toroidal directions.

This idealization, however, breaks down at the plasma edge where both the magnetic topology and various perpendicular transport processes introduce at least a poloidal asymmetry.

We show that the mass flows and radial electric field driven by edge poloidal density asymmetries can be used as a highly effective control mechanism for the edge and thus the global confinement in tokamaks. The underlying physics can be demonstrated

entirely within a simple magnetohydrodynamic equilibrium model with an appropriate flow damping mechanism. As an example, strong dependence of the low to high (LH) transition power threshold on the magnetic topology, an experimental observation still poorly understood, can be easily explained within this framework. Similar arguments also indicate that some of the ITER fueling ports are misplaced from an operational point of view and may lead to higher input power requirements.

## Country or International Organization

Korea, Republic of

## Paper Number

TH/P8-2

**Primary author:** Dr AYDEMIR, Ahmet Y. (National Fusion Research Institute, Daejeon, Korea)

**Co-authors:** Dr PARK, ByoungHo (NFRI); Dr LEE, Hyungho (National Fusion Research Institute); Dr SEOL, JaeChun (National Fusion Research Institute); Dr IN, Yongkyoon (National Fusion Research Institute)

**Presenter:** Dr AYDEMIR, Ahmet Y. (National Fusion Research Institute, Daejeon, Korea)

**Session Classification:** P8 Posters