Contribution ID: 420

Type: Poster

## The Effect of Pressure Anisotropy on Ballooning Modes in Tokamak Plasmas

Thursday 25 October 2018 08:30 (20 minutes)

Edge Localised Modes (ELMs) are thought to be caused by a spectrum of magnetohydrodynamic instabilities, including the ballooning mode. While ballooning modes have been studied extensively both theoretically and experimentally, the focus of the vast majority of this research has been on isotropic plasmas. The prevalence of pressure anisotropy in modern tokamaks thus motivates further study of these modes. This paper presents a numerical analysis of ballooning modes in anisotropic equilibria. The investigation was conducted using the newly-developed codes HELENA+ATF and MISHKA-A, which adds anisotropic physics to equilibria and stability analysis. We have examined the impact of anisotropy on the stability of an n=30 ballooning modes in equilibria with different levels of anisotropy were then calculated using the stability code MISHKA-A. The key finding is that the level of anisotropy had a significant impact on ballooning mode growth rates. For  $T\perp$  >  $T\parallel$ , typical of ICRH heating, the growth rate increases, while for  $T\perp$  <  $T\parallel$ , typical of neutral beam heating, the growth rate configurations to be significant. An important conclusion is the possibility that higher ELM-free performance might be achieved by increasing  $p\parallel/p\perp$  in the pedestal region.

## **Country or International Organization**

Australia

## Paper Number

TH/P5-12

Author: Dr HOLE, Matthew John (Australian National University)

**Co-authors:** Mr JOHNSTON, Alex (Australian National University); Mr BARNES, Ashley (Monash University); Mr HEZABEH, Houman (Australian National University); Dr QU, Zhisong (Australian National University)

Presenter: Dr HOLE, Matthew John (Australian National University)

Session Classification: P5 Posters