

# International Conference on the Management of Naturally Occurring Radioactive Materials (NORM) in Industry

VIRTUAL EVENT

International Conference on  
**Management of Naturally  
Occurring Radioactive  
Material (NORM) in Industry**

19–30 October 2020

#NORM2020



Contribution ID: 18

Type: Poster

## Radioactivity concentrations and exposure rates near a coal power plant in Greece

Coal-fired power plants are the main source of electricity in Greece. Four thermal power stations with more than 4000 MW total installed capacity are located in Northern Greece, producing more than 70% of the country's power requirements.

About 64 Mt of lignite are produced annually by opencast mining in four mines and are transported to the power stations by rail trucks and conveyor belts. Given that, around 15% of the fuel is converted to ash, and assuming 99.9% collection efficiency for the electrostatic precipitators, the amount of fly ash expected to be released annually into the basin's atmosphere is nearly 9.6 kt.

A survey around the Coal Power Plant has been conducted by using portable gamma ray spectrometers, studying an area of 25000m<sup>2</sup> in a grid of 500x500 m in places that were accessible and 1000x1000 at non-accessible places. Two different instruments have been used (a) Saphymo-Stel 1"NaI(Tl), Type S.P.P. 2NF and (b) Identifier, target system electronic gmbh. Direct measurement on the level of 1 m above the ground detected the dose rate in each sampling point.

Thirty six (36) soil samples have been collected and analyzed by gamma-spectrometry in the laboratory for validation of the obtained in-situ measurements. Concentrations of <sup>226</sup>Ra, <sup>232</sup>Th, <sup>40</sup>K and <sup>137</sup>Cs ranged between 20-42 Bq kg<sup>-1</sup>, 13-43 Bq kg<sup>-1</sup>, 260-520 Bq kg<sup>-1</sup> and 20-28 Bq kg<sup>-1</sup> respectively.

The estimated dose rate were calculated by the following equations:

$$D=0.0414 A(40K)+0.623 A(232Th) +0.461 A(226Ra)$$

$$D=0.0414 A(40K)+0.623 A(232Th) +0.461 A(226Ra)+1.72 F(137Cs)$$

where,

D, is the absorbed dose rate in air in nGy h<sup>-1</sup> per Bq kg<sup>-1</sup>

A, the activity concentrations in Bq kg<sup>-1</sup> of each one radionuclide in soil

<sup>137</sup>Cs 1.72 nSv h<sup>-1</sup> per kBq m<sup>-2</sup>

F, the <sup>137</sup>Cs activity concentrations in soil in Bq m<sup>-2</sup>

A coefficient of 0.7 Sv Gy<sup>-1</sup> converts absorbed doses in air to effective dose equivalent

Prevailing winds are weak to moderate, blowing mostly along the NW/SE axis of the basin resulting in higher concentrations at this direction.

The measured external doses are in agreement with the estimated external doses. The observed external doses 30-93 nSv h<sup>-1</sup> are in agreement with the population-weighted average absorbed dose rate in air from terrestrial gamma radiation 57 nGy h<sup>-1</sup> and the reported in Greece average absorbed dose rate in air from terrestrial gamma radiation 42 nGy h<sup>-1</sup>.

The results can be used as a reference point for radiological mapping of the studied area.

**Primary authors:** IOANNIDOU, ALEXANDRA; Ms BETSOU, Chrysoula (Aristotle University of Thessaloniki, Physics Department)

**Presenter:** IOANNIDOU, ALEXANDRA

**Session Classification:** Session IV - Characterization in Industrial Facilities and in the Environment

**Track Classification:** NORM Characterization, Measurement, Decontamination