

# Vessel Forces from a Vertical Displacement Event in ITER

Wednesday, May 12, 2021 12:10 PM (20 minutes)

1. Introduction : Disruptions are one of the major concerns in ITER and other future tokamaks [1]. A particularly troublesome type of disruption is a vertical displacement event (VDE) where control of the vertical position of the plasma column is lost. In addition to heat, particle flux, and energetic electrons impacting the first wall, significant electromagnetic loads will arise. For realistic modelling of a VDE disruption, a detailed 3D model of the disrupting plasma and an accurate description of the conducting structures surrounding the plasma is required. The structure affects the plasma evolution itself and the plasma acts as a source of currents and fields which produce the electromagnetic loads. Most of the VDE modeling work to date has used the axisymmetric evolving equilibrium codes TSC [2], DINA [3], and CarMa0NL[4] to describe the disrupting plasma. This paper describes our efforts to extend this analysis by using the fully 3D MHD codes M3D-C1 [5], NIMROD [6], and JOREK [7]. We describe our efforts in benchmarking these 3 codes on VDE relevant calculations, validation with some experimental data, and projection to ITER vessel and plasma conditions. Attention is given to the role of the currents shared by the plasma and the structure (halo currents).
2. 2D and 3D benchmarking of M3D-C1, NIMROD, JOREK in simplified geometry We have undertaken two VDE code benchmarking exercises for the 3 MHD codes M3D-C1, NIMROD, and JOREK. The first was in 2D (axisymmetric) [8] and we obtained excellent agreement for the linear and early nonlinear growth rates and for the motion of the magnetic axis, induced wall currents, and distribution of the halo current. The second benchmark, currently in progress, is fully 3D, beginning at the time the plasma first makes contact with the vessel.
3. Simulation of the sideways force in JET: M3D-C1 simulations of asymmetric vertical displacement events (AVDEs) in JET are in progress. The simulations were compared with previous M3D simulations [9] and with the JET 2011-2016 ILW disruption database. The computed and experimentally measured Noll approximations to the sideways force are in reasonable agreement. An important feature of the simulations and the data is that the sideways wall force is small when the current quench time is less than the wall penetration time.
4. 2D benchmark of M3D-C1 and CarMa0NL for ITER geometry with no halo currents We performed a separate nonlinear but axisymmetric VDE benchmark between M3D-C1 and CarMa0NL using a uniform thickness resistive wall located at the position of the ITER first wall. A 6 cm thick wall with resistivity of  $7.4E-7$  ohm-m gives a time constant of 235 ms for the wall current to develop when a constant loop voltage is applied (no plasma). Modeling of both a "hot" and "cold" VDE with the 2 codes, both without halo currents present, gave excellent agreement. [10]
5. The effect of Halo Currents on the vertical force: We have performed extensive axisymmetric studies with M3D-C1 of the effect of different assumptions regarding the temperature distribution of the halo region during an axisymmetric VDE in ITER geometry with the wall model described in #4 above. These studies use spatially and time varying parallel and isotropic thermal conductivities to model the thermal quench which then caused the current quench with differing amounts of halo current. We found the result that when a larger force from the halo current is present, it is at least partially offset by a smaller force due to the induced toroidal currents [11]. Slower current quenches generally lead to larger vertical forces, but the presence of a halo region can modify this scaling. Also, sheath-type boundary conditions are found to slow the evolution, increasing impulse in NIMROD computations [12].
6. 3D VDE simulations in ITER geometry: Building on these successful code benchmarks and axisymmetric parameter studies in ITER geometry, we have performed several 3D VDE simulations with M3D-C1 using the ITER vessel model described in 4.




Figure 1:

Currents and fields from these simulations are input to the CARRIDI code for detailed force calculations. Our initial result is that the intrinsically 3D “sideways force” is much smaller than the vertical force, largely because the kinking motion of the plasma occurs on a time scale much faster than the vessel time constant. This is largely consistent with an earlier scaling obtained with the M3D code. [13]

7. Summary and Conclusions: For the first time, we have applied massively parallel implicit 3D magnetohydrodynamics codes to predict the wall forces produced during ITER disruptions using realistic values for the wall resistivity and time constant. Extensive code verification and validation has been performed. We are in the process of refining the wall model to make it more realistic, and in developing a tighter coupling of M3D-C1 with CARRIDI in order to refine our present results. Acknowledgments: This work was supported by US DOE Contracts No. DE-AC02-09CH11466, and No. DE-SC0018001 the SciDAC Center for Tokamak Transient Simulations, EUROfusion Researcher Fellowship programme

under the task agreement WP19-335 20-ERG-DIFFER/Krebs, EUROfusion grant No. 633053, F4E contract F4E-OPE-0891), and by Italian MIUR under PRIN grant 20177BZMAH. Disclaimer ITER is the Nuclear Facility INB no. 174. The views and opinions expressed herein do not necessarily reflect those of the ITER Organization. This publication is provided for scientific purposes only. Its content should not be considered as commitments from the ITER Organization as a nuclear operator in the frame of the licensing process. References:

8. T. Schioler, C. Bachmann, G. Mazzone, et al, Fusion Engin. Design, 86, 11, pp. 1963-1966, (2011).
9. S. C. Jardin, N. Pomphrey, and J. Delucia, J. Comp. Phys., 66, pp. 481–507 (1986).
10. R. Khayrutdinov and V. Lukash, J, Comp. Phys., 109, pp. 193–201 (1993)
11. F. Villone et al., Plasma Phys. Control. Fusion 55 (2013) 09500
12. S. Jardin, N. Ferraro, J. Breslau et al, Comput. Sci. & Disc., 5 p 014002 (2012)
13. C. Sovinec, A. Glasser, T. Gianakon, et al, J. Comp. Phys, 195 pp. 355386 (2004)
14. M. Hoelzl, P. Merkel, G. Huysmans, et al. JPCS, 401, 012010 (2012)
15. I. Krebs, F. J. Artola, C. R. Sovinec, et al., “Axisymmetric simulations of vertical displacement events in tokamaks: A benchmark of M3D-C1, NIMROD and JOREK”, submitted to Phys. Plasmas (2020)
16. H. Strauss, S. Jachmich, E. Joffrin, V. Riccardo, J. Breslau, R. Paccagnella, and JET contributors, Physics of Plasmas 24, 102512 (2017)
17. S. Jardin, F. Villone, C. Clauser, N. Ferraro, N. Isernia, G. Rubinacci, S. Ventre, “ITER Disruption Simulations with Realistic Plasma and Conductors Modelling”, Proceedings of EPS, (2019)
18. C. F. Clauser, S. C. Jardin, and N. M. Ferraro, Nuclear Fusion 59 126037 (2019)
19. K. J. Bunkers, PhD Dissertation, University of Wisconsin-Madison (2019)
20. H. Strauss, R. Paccagnella, and J. Breslau, Phys. Plasmas 17, 082505 (2010)

## Country or International Organization

United States

## Affiliation

Princeton Plasma Physics Laboratory

**Primary author:** Dr JARDIN, Stephen (Princeton Plasma Physics Laboratory)

**Co-authors:** Dr CLAUSER, C.; Dr FERRARO, N.; Dr KREBS, I.; Dr ARTOLA, F. J. (ITER Organization, Route de Vinnon sur Verdon, 13067 St. Paul Lez Durance, Cedex, France); Dr BUNKERS, K. (University of Wisconsin-Madison, Madison WI 53706-1609, USA); Prof. SOVINEC, C. (University of Wisconsin-Madison); Dr HOELZL, M. (Max Plank Institute for Plasma Physics); Prof. VILLONE, F. (Consorzio CREATE, DIETI); Dr RUBINACCI, G. (Consorzio CREATE, DIETI); Mr ISERNIA, N. (Consorzio CREATE, DIETI); Dr STRAUSS, H. (HRS Fusion)

**Presenter:** Dr JARDIN, Stephen (Princeton Plasma Physics Laboratory)

**Session Classification:** P3 Posters 3

**Track Classification:** Magnetic Fusion Theory and Modelling