

Spontaneous and triggered abrupt and non-local reduction of electron heat and density fluxes and ITB formation in T-10 tokamak plasmas with ECRH/ECCD

Wednesday 12 May 2021 17:40 (20 minutes)

During non-local ("global") L-H transitions found in various regimes of JET and JT-60U tokamaks earlier [1-3], the rise of $T_{e,i}$ and n_e starts simultaneously in the spatial zone $\approx 0.3 < r/a < 1$, while heat and density fluxes fall simultaneously in the same region. At the ITB-events in JT-60U and T-10 (circular tokamak with a limiter ≈ 30 cm, $R=150$ cm, $B_T=2.5$ T), heat and density fluxes fall in a narrower internal spatial zone within ≈ 40 -50% of the minor radius, see detail in [3-5].

The present report describes the new type of L-H transitions in plasmas with W-limiter and Li-coating called "semi-global". T_e starts to rise simultaneously at $0.2 < r/a < 0.6$ similar to its behavior during an ITB-event in JT-60U and T-10. The density rises in a wider zone $\approx 0.3 < r/a < 1$ similar to a non-local L-H transition. The electron heat flux abruptly reduces in the spatial zone $0.2 < r/a < \approx 1$. The gradual formation of ITB occurs after transition. The rise of T_e in the internal part of plasmas was mentioned at L-H transitions in circular tokamaks with the limiter TUMAN-3, JIPP T-IIU and TEXT-U [6-8] (light cooling of the periphery, like in our cases, was mentioned also). The formation of ITB measured with Thomson scattering was reported in [6]. Nevertheless, analysis of the non-locality of the electron heat flux jump was not done in [6-8].

The one spontaneous semi-global L-H transition was briefly described by us at EC-2018 workshop [9]. The spontaneous transitions (including dithering one) are observed at simultaneous co+contr ECCD by 2 gyrotrons only (total power 1.4-1.5 MW). The presence of dithering transitions (up to 5 H-mode phases with 5-10 ms length) means that the power is near critical value. The figure 1(a) shows the typical evolution of $T_e(r,t)$, line-averaged central density and energy content W at spontaneous and triggered semi-global transitions. In this particular case, neon puffing starts 5 ms before transition ($PECRH=0.85$ MW, see experiments [10]) and puffing starts 15 ms before transition in the regime with higher density and EC-power [11]. The plasma-wall interaction falls (D-beta level drops), T_e starts to rise at $0.2 < r/a < 0.6$ together with the rise of density and W . The density at the edge increases 30% faster compare with that of in the bulk plasmas. The transition triggered by neon puffing is not a transition to RI mode since the level of radiation losses is small enough (below $\sim 25\%$) and just starts to rise before the transition. Figure 1 (b) represents the typical profile of the electron heat flux $\Delta(G_{Te})$ at the transitions in all cases. In this case the transition at $PEC=1.4$ MW and co+co ECCD, was caused by spontaneous drop of Li-containing flake (jump of flux is constant in space up to the edge if one believe to ECE data at the periphery). In a contrast with [1-3], the rise of density is important at calculations of $\Delta(G_{Te})$ profile at transitions on T-10 by simple expression [1-5]: $\Delta(G_{Te}(r))S(r) = 3/2 \{ \text{volume integral of } \Delta[d(nTe)/dt] \}$, $S(r)$ – enclosed surface and $\Delta[d(nTe)/dt]$ is the jump of derivative at the time of transition with the evolution of $\Delta(n(r,t))$ calculated from data derived by 16 channels interferometer. The transport gradually varies in space and time after transition and leads to the formation of ITB as one can see on the figure 1(b) (typical case). The similar case in the plasmas with co+contr ECCD has been described recently in [12].

The authors of the paper [13] show that the rise of central T_e at the injection of small carbon pellets at ASDEX-U is explained by strong stabilization of the trapped electron mode turbulence due to the measured fast significant flattening of the density profile in the internal part of plasma column. This explanation is not valid in our cases since the density just starts to rise gradually after transition (even at the drop of flake the density varies abruptly at the edge only).

The paper describes the new type of transition called "semi-global" L-H transition. T_e starts to rise simultaneously at $0.2 < r/a < 0.6$ similar to its behavior during an ITB-event in JT-60U and T-10. The density rises in a wider zone $\approx 0.3 < r/a < 1$ similar to a global L-H transitions at JET and JT-60U and the electron diffusivity falls in the same zone. The electron heat flux abruptly reduces in the spatial zone $0.2 < r/a < \approx 1$. The spontaneous transitions (including dithering one) are observed at simultaneous co+contr ECCD by 2 gyrotrons only. New triggers of the transitions are neon gas puffing and spontaneous drop of the Li-containing flakes in various regimes of ECCD and EC-power (observed at $n_{line\ av}(0) < 3 \cdot 10^{19}/m^3$, $B_T=2.3$ -2.5 T, $I_p=200$ -250 kA). By our up to date knowledge, transitions with the non-local reduction of heat flux has never been reported in the tokamaks with limiter. An abrupt increase of energy confinement time τ_E at the moment of the transition is equal to 10-20% (small value of H-factor is typical at circular tokamaks). The H mode continues up to the double value of τ_E . The accumulation of impurities is absent. The part of T_e rise can be explained by the abrupt reduction of the convective electron heat flux $2.5T_eG_n$ due to the decay of electron density flux G_n at the time of transition.

The authors thank Drs A.Ya Kislov, D.A. Kislov, N.A. Kirneva and many other members of the T-10 team for fruitful discussions and fine collaboration. The work was supported by ROSATOM Corporation.

References

1. Neudatchin S V, Cordey J G and Muir D J 20th EPS Conf. on Control. Fus. and Plasma Phys. (Lisboa) vol. I (Geneva : EPS), p 83 (1993)
2. Neudatchin S V, Takizuka T, Shirai H et al., Japan J. Appl. Phys. 35 3595 (1996)
3. Neudatchin S. V., Takizuka T., et al., Plasma Phys. Control. Fusion 44 A383-389 (2002)
4. Neudatchin S V., Inagaki S, Itoh K et al. 2004 J. Pl. and Fus. Res. Series 6 134
5. Neudatchin S.V, Shelukhin D.A., Mustafin N.A. 2017 J. Phys.: Conf. Ser. 907 012015
6. Andrejko M V, et al 1992 IAEA_PPCNFR_Wuerzburg_v1_p485 IAEA-CN-56/A-7-11
7. Watari T., Kumazawa R. et al 1990 Nucl. Fusion 30 1197
8. Roberts D. R. et al 1996 Plasma Phys. Control. Fusion 38 1117
9. Borshegovskiy A, Neudatchin S, Pimenov I et al 2019 EPJ Web of Conf. 203 02004
10. Kasyanova N.V., Rasumova K.A. et al, . 2018, in Procs. of 45th EPS Conf. on Pl. Ph, Prague, ECA, Vol 42A, P4. 1106
11. Kirneva N.A. et al, 2018, 45th EPS Conf. on Pl. Ph., Prague, ECA, Vol 42A, P4. 1081
12. Neudatchin S. V., Pimenov I. S, et al., 2019, J. of Ph.: Conf. Ser. 1383 012005
13. Angioni C. et al 2019 Nucl. Fusion 59 106007

Indico rendering error

Could not include image: [429] Error fetching image

Indico rendering error

Could not include image: [429] Error fetching image

Affiliation

NRC Kurchatov Institute

Country or International Organization

Russia

Author: Dr NEUDATCHIN, Sergey (NRC Kurchatov Institute)

Co-authors: Mr KYUANOV, Aleksey (NRC Kurchatov Institute); Dr BORSCHEGOVSKIY, Alexey (NRC Kurchatov Institute); Mr PIMENOV, Igor (NRC Kurchatov Institute)

Presenter: Dr NEUDATCHIN, Sergey (NRC Kurchatov Institute)

Session Classification: P4 Posters 4

Track Classification: Magnetic Fusion Experiments