Contribution ID: 132

Type: Invited

Mitigation of runaway electron heat loads by deuterium SPI injection and kink activity

Runaway electrons (RE) are a major threat for a reliable operation of future tokamaks including ITER. Avoiding or dissipating them is therefore essential. Shattered Pellet Injection (SPI) is the disruption mitigation and RE avoidance method currently planned for ITER [1]. However, if this first line of defence is not efficient enough to prevent the formation of REs, SPI must also be able to dissipate a fully formed RE beam. Recent experiments on JET [2] and DIII-D [3,4] using deuterium SPI showed complete dissipation of RE beams up to 1.2 MA. No measurable localized impact and damage on plasma facing components is observed in the JET cases. The runaway current is converted into ohmic current which then decays as a "standard" plasma current quench. Conversely, past experiments using Massive Gas Injection (MGI) or SPI had shown that beam mitigation using neon, argon or krypton was at best incomplete [5,6]. The RE current could be decreased in selected situations (low density companion plasma) but the final RE impact on the wall could not be avoided, leading to localized melting with currents as low as 200 kA [7].

The mechanisms leading to complete RE loss using deuterium SPI are twofold: a large MHD instability and a low-impurity content companion plasma. The companion plasma following D2 SPI is purged out of its remaining high-Z impurities, with electron temperature less than a few eV and electron density lower than 10^{18} m⁻³. Current is allowed to grow again due to the voltage from the central solenoid, ultimately leading to a low-q MHD instability, most likely a kink [7]. MHD simulations with MARS-F and JOREK show that such a large amplitude kink leads to enhanced RE losses and increased wetted area. The reason why the instability is stronger with D2 SPI is not yet fully understood, but may be related to a more hollow RE current profile following the deuterium SPI. Such a profile can exhibit current-driven kink instability at any integer qedge. Once the instability has dissipated the RE plateau, no magnetic energy is converted back to RE kinetic energy thanks to the low impurity content: the current quench is slow enough and its bound electron density low enough to prevent re-avalanching of RE. The conversion mechanism is much stronger in high-Z SPI mitigation cases compared to D2 SPI cases, thus explaining why high-Z runaway mitigation is incomplete.

These experiments and simulations show that D2 SPI is a promising RE mitigation mechanism for ITER in case a RE beam appears. Simulations are underway to quantify the degree of high-Z purging which is needed to achieve a sufficiently low conversion rate from magnetic to kinetic energy following the MHD instability. Acknowledgments: This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 and 2019-2020 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission. Work supported by US DOE under DE-FC02-04ER54698 and DE-SC0020299.

Member State or International Organization

France

Affiliation

CEA, IRFM, F-13108 Saint-Paul-les-Durance, France

Primary authors: REUX, Cedric (CEA, IRFM, F-13108 Saint Paul-lez-Durance, France.); PAZ-SOLDAN, Carlos (General Atomics); HOLLMANN, Eric M. (University of California San Diego); EIDIETIS, Nicholas (General Atomics); LEHNEN, Michael (ITER Organization); JACHMICH, Stefan (ITER Organization); JOFFRIN, Emmanuel (CEA); Dr LOMAS, Peter (UKAEA/CCFE); Dr RIMINI, Fernanda (CCFE, Culham Science Centre, Abingdon, Oxon, OX14 3DB, United Kingdom of Great Britain and Northern Ireland); BANDARU, Vinodh Kumar (Max-Planck-Institute for Plasma Physics, Garching); BAYLOR, Larry R. (Oak Ridge National Laboratory); Dr IGOR, Bykov (University of California-San Diego, 9500 Gilman Dr., La Jolla, CA 92093-0417, United States of America); Mr CALACCI, Luca (Università di Roma Tor Vergata, Via del Politecnico 1, Roma, Italy); CAUSA, Federica (ENEA C. R. Frascati); CARNEVALE, daniele (Universita' Roma Tor Vergata, Dipartimento di Ing. Civile ed Ing. Infor-

matica); Dr COFFEY, Ivor (UKAEA/CCFE); Mr CRAVEN, Douglas (UKAEA/CCFE); Mr DAL MOLIN, Andrea (University Milano-Bicocca, Piazza della Scienza 3, 20126 Milano, Italy); DE TOMMASI, Gianmaria (Università degli Studi di Napoli Federico II); DU, xiaodi (General Atomics); FICKER, Ondrej (Institute of Plasma Physics of the Czech Academy of Sciences); GEBHART, Trey (Oak Ridge National Laboratory); GERASIMOV, Sergei (CCFE); Dr GIACOMELLI, Luca (Istituto per la Scienza e Tecnologia dei Plasmi, ISTP- CNR, via R. Cozzi 53, 20125 Milano, Italy); HERFINDAL, Jeffrey (UsOakRidge); HOELZL, Matthias (Max Planck Institute for Plasma Physics); HUBER, Alexander (Forschungszentrum Jülich GmbH, Institut für Energie- und Klimaforschung -Plasmaphysik); Dr LASNIER, Charlie (Lawrence Livermore National Laboratory); LIU, Yueqiang (General Atomics, PO Box 85608, San Diego, CA 92186-5608, USA); LVOVSKIY, Andrey (Oak Ridge Associated Universities); LOWRY, Christopher (European Commission); Dr MACUSOVA, Eva (Institute of Plasma Physics of the CAS); Prof. MAN-ZANARES, Ana (Laboratorio Nacional de Fusión, CIEMAT); MCLEAN, Adam (Lawrence Livermore National Laboratory); NOCENTE, Massimo (Dipartimento di Fisica, Università di Milano-Bicocca); Mr PANONTIN, Enrico (University Milano-Bicocca, Piazza della Scienza 3, 20126 Milano, Italy); PAUTASSO, Gabriella; Dr PEACOCK, Alan (UKAEA/CCFE); PLYUSNIN, Vladislav V (Instituto de Plasmas e Fusão Nuclear, Associação EURATOM-IST, Instituto Superior Tecnico); RUDAKOV, D. L. (University of California San Diego); SHIRAKI, Daisuke (Oak Ridge National Laboratory); Dr SILBURN, Scott (UKAEA/CCFE); Dr SOMMARIVA, Cristian (Ecole Polytechnique Fédérale de Lausanne (EPFL), Swiss Plasma Center (SPC), CH-1015 Lausanne, Switzerland); SOZZI, Carlo (Istituto per la Scienza e Tecnologia dei Plasmi ISTP-CNR Milano Italy); SRIDHAR, Sundaresan (IRFM-CEA Cadarache); Dr SWEENEY, Ryan (MIT PSFC); Dr TINGUELY, Alex (MIT PSFC)

Presenter: REUX, Cedric (CEA, IRFM, F-13108 Saint Paul-lez-Durance, France.)

Track Classification: Mitigation