

# Evaluation of extreme ultraviolet spectral models for mid-charged tungsten ions with LHD experiments

Monday, 15 May 2023 10:40 (20 minutes)

The behavior of tungsten impurity in fusion plasmas is one of the important issues to be studied to achieve high-temperature plasmas for fusion reactions since tungsten used as plasma-facing material is sputtered by plasma particles and is expected to reduce electron temperature due to large radiation power. Many studies have been done to examine atomic properties and spectral models for tungsten ions, however, tungsten ions with 4f open subshells are not well studied yet due to the complexity of the atomic structure. Extreme ultraviolet (EUV) spectra of tungsten ions have been measured in various fusion devices and electron beam ion traps (e.g. [1,2]), and examined by comparison of theoretically calculated spectra by collisional-radiative (CR) models (e.g. [3,4]). The unresolved transition array (UTA) measured at 4.5-7nm wavelength region for plasma with electron temperature  $\sim 1$ keV is produced by numerous overlapped 4d-4f and 4p-4d transitions of tungsten ions. The wide two-peak feature of the UTA profile is not fully understood yet, even though recombination processes are included in the CR models [4,5] for  $Wq^+$  with  $q=25-39$ . On the other hand, the peaks measured at 2-4nm are well understood and are produced by many  $n=4-5$  and  $n=4-6$  transitions of  $Wq^+$  with  $q=22-30$ . They are useful to estimate charge state distributions [4,5]. For ions with  $q<22$ , no peaks are found in this region and we need some identifier for such lower charged ions. We extend our study to EUV spectra at 10-30nm where  $n=5-5$  transitions are found for mid-charged tungsten ions by CR model calculations. We have performed plasma experiments to measure tungsten spectra by pellet injection with Large Helical Device (LHD) for wide wavelength regions. Measured spectra at 10-30nm can be used to evaluate calculated spectra by CR models for mid-charged tungsten ions. Details of the comparison will be presented at the conference.

- [1] R. Neu et al., Plasma Phys. Control. Fusion 38, A165 (1996)
- [2] H. A. Sakaue et al., AIP Conf. Proc. 1438, 91 (2012)
- [3] T. Putterich, R. Neu, R. Dux et al., Plasma Phys. Control. Fusion 50, 085016. (2008)
- [4] I. Murakami, H. A. Sakaue, C. Suzuki et al., Nucl. Fusion 55, 093016 (2015)
- [5] I. Murakami, D. Kato, T. Oishi et al., Nucl. Mater. Energy 26, 100923 (2021)

## Presenting Author

Izumi Murakami

## Presenting Author Affiliation

National Institute for Fusion Science

## Presenting Author Gender

Female

## Country

Japan

## Presenting Author Email Address

murakami.izumi@nifs.ac.jp

**Primary author:** Prof. MURAKAMI, Izumi (National Institute for Fusion Science)

**Co-authors:** Dr SASAKI, Akira (National Institute for Quantum Science and Technology); Dr KATO, Daiji (National Institute for Fusion Science); Dr SAKAUE, Hiroyuki A. (National Institute for Fusion Science); Dr GOTO, Motoshi (National Institute for Fusion Science); Dr OISHI, Tetsutarou (Tohoku University); Dr KAWATE, Tomoko (National Institute for Fusion Science); Dr KAWAMOTO, Yasuko (National Institute for Fusion Science)

**Presenter:** Prof. MURAKAMI, Izumi (National Institute for Fusion Science)

**Session Classification:** Magnetic-Confinement Fusion Plasmas

**Track Classification:** Magnetic-Confinement Fusion Plasmas