

Simulation of Doppler-free Spectra using the Collisional Radiative Model

Wednesday, 17 May 2023 11:40 (20 minutes)

Saturated absorption spectroscopy is a tool that can be used to suppress the Doppler broadening of observed atomic and molecular transition lines in order to measure their precise wavelengths. Obtaining a saturated absorption condition by laser excitation is an essential technique for use in saturated absorption spectroscopy. We are introducing the laser excitation process into the collisional radiative model of hydrogen atoms to uncover how much saturation can be achieved under realistic plasma conditions and laser power density. Results show that the simulated spectra were able to successfully model Lamb dips and peaks utilizing this method, with the simulated plasma and laser parameters showing good agreement with the ones used in the experiment. This model has additionally helped to give further insight into how plasma parameters can affect the spectral characteristics of Lamb dips and peaks by noting the dependence of the simulated spectral saturation on these parameters.

Presenting Author

Joseph John Simons

Presenting Author Affiliation

Department of Fusion Science, The Graduate University for Advanced Studies, SOKENDAI

Presenting Author Gender

Male

Country

Japan

Presenting Author Email Address

josephjohn.simons@nifs.ac.jp

Primary author: SIMONS, Joseph John (Department of Fusion Science, The Graduate University for Advanced Studies, SOKENDAI)

Co-author: GOTO, Motoshi (National Institute for Fusion Science)

Presenter: SIMONS, Joseph John (Department of Fusion Science, The Graduate University for Advanced Studies, SOKENDAI)

Session Classification: Fundamental Data and Modelling

Track Classification: Fundamental Data and Modelling