

Unstable beta-induced ion temperature gradient (BTG) eigenmodes in JET plasmas with ITBs and elevated monotonic q-profiles.

Monday, 6 December 2021 14:10 (30 minutes)

Abstract:

JET deuterium experiments in an advanced tokamak scenario with an internal transport barrier (ITB) exhibit unstable electromagnetic (EM) perturbations in the sub-TAE frequency range. In JET pulse number (JPN) 92054, a high-beta plasma ($\beta_N = \beta_T B_T a / I_P \sim 4.38 [\% Tm/MA]$) with high power neutral beam injection (NBI), $P_{NBI} = 25.1 MW$, contained EM perturbations identified as beta-induced ion temperature gradient (BTG) eigenmodes and not beta-induced Alfvén eigenmodes (BAE) nor beta-induced Alfvén acoustic eigenmodes (BAAE) which are often destabilised in similar plasma conditions. The EM perturbations are localised near the $q = 2$ magnetic surface related to the ITB, and their frequency correlates well with the BTG characteristic frequency (ion diamagnetic frequency, ω_i^*) and the thermal ion temperature gradient (∇T_i). BTG modes are the most unstable modes due to the high thermal ion temperature gradient in the ITB, high thermal ion temperature compared to thermal electron temperature ($T_i/T_e > 1$), and a high ion beta. Three well-defined conditions for BTG modes to exist, defined by BTG analytical theory [1], are fulfilled in JPN 92054: (1) a positive relative ion temperature gradient, (2) ion beta higher than a critical value, and (3) a low magnetic shear. BTG theory also predicts a mode location in the vicinity of a rational magnetic surface, a frequency scaling with ω_i^* , and a coupling between Alfvén and drift waves. We have performed linear gyrokinetic simulations with validated plasma profiles and equilibrium, and find a mode with features resembling those of the experimental and theoretical BTG modes; specifically the mode is kinetically driven by thermal ions, is localised near the $q = 2$ magnetic surface, has a dominant Alfvénic polarisation, and its frequency scales with ω_i^* dependent on the toroidal mode number (n). Parts of this work have been reported in [2].

BTG modes are also observed in more recent JET plasmas during energetic particle scenario experiments aimed at studying alpha-particle driven AEs, performed in JET 2019/2020 deuterium campaigns. Reflectometer diagnostic data confirm that the mode location is around the $q = 2$ magnetic surface. We also present evidence for a systematic correlation between the BTG mode stability and the neutron rate roll-over (i.e. $d(R_{NT})/dt$ transiting from positive to negative).

References:

- [1] A. B. Mikhailovskii and S. E. Sharapov. Beta-induced Temperature-gradient Eigenmodes in Tokamaks. Kinetic Theory. JET Joint Undertaking Reports, JET-P(98)12:1-16, 1998.
- [2] N. Fil, et al. Interpretation of electromagnetic modes in the sub-TAE frequency range in JET plasmas with elevated monotonic q-profiles. Physics of Plasmas, Accepted, 2021.

Acknowledgments:

This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 and 2019-2020 under grant agreement No 633053 and from the RCUK [grant number EP/T012250/1]. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

This work was supported by U.S. Department of Energy (DOE) through DEFG02-99ER54563, DE-AC05-00OR22725 and DE-AC02-05CH11231).

Speaker's Affiliation

UKAEA, CCFE, Abingdon.

Member State or IGO

United Kingdom

Primary author: Dr FIL, Nicolas (UKAEA, CCFE)

Co-authors: SHARAPOV, Sergei (CCFE); Dr FITZGERALD, Michael (UKAEA); CHOI, GYUNGJIN; LIN, Zhi-hong (UC Irvine); TINGUELY, Roy (MIT PSFC); Dr PUGLIA, Paulo (EPFL); MCCLEMENTS, Ken (CCFE); OLIVER, James (UKAEA, CCFE); DUMONT, Remi (CEA, IRFM); Dr KEELING, David (UKAEA); Dr ANTONIO .C.A. , Figueiredo (IPFN); PORKOLAB, Miklos (MIT); MAILLOUX, J (UKAEA); JOFFRIN, Emmanuel (CEA); AND JET CONTRIBUTORS

Presenter: Dr FIL, Nicolas (UKAEA, CCFE)

Session Classification: Effects of Energetic Particles in Magnetic Confinement Fusion Devices

Track Classification: Effects of Energetic Particles in Magnetic Confinement Fusion Devices