



Assessment of Occupational Exposure from Patient Treated with ¹³¹I for Thyroid Cancer

Y. LAHFI and O. ANJAK

Atomic Energy Commission of SYRIA, Damascus, Syrian Arab Republic

y.lahfi@aec.org.sy

Poster ID: P-S8-141

1. INTRODUCTION

- Oral administration of large doses of radioiodine ¹³¹I has been a commonly accepted procedure for treatment of benign and malignant conditions of the thyroid since 1940s [1].
- Several research papers and guidelines were focused on the external dose rate from patient treated by radioiodine ¹³¹I in discussing the release criteria from hospital [2] or assessing the related public or occupational exposures [3,4].
- The goal of the present work is to provide a simple equation to assess the medical staff's occupational exposure nearby a radioiodine ¹³¹I administrated patient.

2. MATERIALS AND METHODS

- A randomized sample of 86 patients administrated with ¹³¹I for thyroid radioiodine therapy at AL-BAYROUNI University Hospital was monitored.
- The patients' administered activities ranged from 3060 to 5650 MBq.
- A calibrated universal survey meter type RADOS-200 was used to measure the external radiation dose rate from the administrated patients.
- The Measurements were effectuated post radioiodine dose administration at a distance of 1 meter from the effective point of measurement and standing point of the patient at the thorax level.
- The patient's sample was divided into two groups according to the two administrated activity ranges. The patients' age, height, and weight were also recorded.
- Tables 1 and 2 represent the description of the patient groups and the related anthropometric measurements.

TABLE 1. The description of the patient groups

Patient Group	Administered Activity Range Mbq (mCi)	Number of Patients	
		Male	Female
Group A	2960 - 3700 (80 - 100)	10	47
Group B	3737- 5550 (101-150)	5	24

TABLE 2. Anthropometric measurements of patient groups

Patient Group	Group A		Group B	
	Male	Female	Male	Female
	Average ± SD	Average ± SD	Average ± SD	Average ± SD
Age (Year)	42.44±11.11	42.47±13.04	45.60±14.35	38.88±11.64
Length (m)	1.75±0.05	1.60±0.07	1.74±0.05	1.60±0.06
Weight (kg)	99.22±21.45	75.36±13.63	90.40±30.52	75.46±14.60
BMI (kg/m ²)	31.98±5.84	29.42±5.26	29.73±9.74	29.30±6.32

3. RESULTS

- The mean values of the measured patient external radiation dose rates and theirs normalized dose rate values in terms of administered activity for patient groups are presented in Tables 3 and 4.
- The calculated mean values were within the range of 50.3-145 and 75.53-209 μSv.h⁻¹ respectively.
- The mean patient external dose rate in Group B was relatively higher by ~18% than in Group A since the administrated activity in Group B was higher than in Group A, however, the differences in of the external dose rate between female and male was insignificance.

TABLE 3. The external dose rate (μSv.h⁻¹) at a distance of 1 meter from patient post radioiodine ¹³¹I administration (Mean ± SD).

Patient Group	Average external dose rate at 1 meter (μSv.h ⁻¹)		
	Male	Female	Male & Female
Group A	99.37±14.82	104.36±20.45	103.55±19.62
Group B	113.17±35.28	125.12±29.91	123.06±30.56

TABLE 4. The normalized dose rate in μSv.h⁻¹.Mbq⁻¹ at a distance of 1 meter from patient and post radioiodine ¹³¹I administration (Mean ± SD).

Patient Group	Normalized dose rate at 1 meter in μSv.h ⁻¹ .Mbq ⁻¹ (μSv.h ⁻¹ .mCi ⁻¹)		
	Male	Female	Male & Female
Group A	0.031±0.005 (1.15±0.18)	0.031±0.006 (1.16±0.22)	0.031±0.006 (1.16±0.21)
Group B	0.030±0.009 (1.11±0.35)	0.030±0.005 (1.12±0.19)	0.030±0.006 (1.12±0.22)

4. DISCUSSIONS

- The radiation exposure from an administrated patient to nearby persons is related primarily to the distance from him, to the exposure duration, and to the administered radioiodine activity.
- The maximum external dose rate measured from the monitored patient sample was 209 μSv.h⁻¹ and the mean normalized value of the external dose rate was 0.031±006 μSv.h⁻¹.Mbq⁻¹ with a coefficient of variation of 18.7%.
- By applying the inverse square law, the average worker's effective dose nearby a patient administrated with an activity A (Mbq) of radioiodine ¹³¹I at a distance d (m) and for a time duration of T (Hour) can be estimated by the following equation:

$$E(\mu Sv) = 0.031 * T * \frac{1}{d^2} * A$$

- For of a realistic exposure scenario simulating the presence nearby an administrated patient with 3700 Mbq of radioiodine ¹³¹I for 5 minutes at a distance of 50 cm and for 10 minutes at a distance of 1 meter, the estimated worker's effective dose will be 57.35 μSv.
- As the average annual monitored effective dose for exposed workers in a nuclear medicine is ranged from 3 to 5 mSv [5], the proposed equation could be a useful tool to estimate instantly the medical staff's occupational or emergency exposure and consequently to maintain the conformity with radiation protection principles.

5. CONCLUSIONS

- The radiation dose nearby a sample of 86 patients administrated with ¹³¹I for radioactive iodine therapy has been assessed. The average normalized patient external dose rate at 1 meter in term of radioiodine activity was 0.031±006 μSv.h⁻¹.Mbq⁻¹.
- An equation based on the normalized external dose rate was proposed to estimate the average worker's effective dose from an administrated patient. It could be a simple way to estimate instantly the resulted occupational or emergency exposure and to maintain the conformity with radiation protection principles.

6. ACKNOWLEDGEMENTS

The authors express appreciation to Dr. Ibrahim OTHMAN, General Director of AECS and to Dr. Mohammad Said Al-MASRI, Head of Protection and Safety Department (AECS) for their continuous support.

7. REFERENCES

1. ROSS, I., Thyroid Cancer in Clinical Practice, Springer, (2007).
2. Grigsby P, Siegel B, et al, Radiation exposure from outpatient radioactive iodine (¹³¹I) therapy for thyroid carcinoma. JAMA; 283, (2000) 2272-4.
3. Stenstad L.I., Pedersen G.A., Landmark A.D., Brattheim B., Nuclear radiation dose to the surroundings from patients who are undergoing nuclear medicine examinations. Radiography Open, 1 (2014) 10-18.
4. Sattari A., Dadashzadeh S., Nasiroghli G., Firoozabadi H., Radiation dose to the nuclear medicine nurses. Iran. J. Radiat. Res., 2 (2004), 59-62.
5. IAEA, Training Material on Radiation Protection in Nuclear Medicine (2017), https://rpop.iaea.org/RPOP/RPoP/Content/Documents/TrainingNuclearMedicine/Lectures/RPNM_Par05_worker_WEB.ppt