

Modeling the effect of drifts on HL-2M snowflake in-out divertor heat loading by SOLPS-ITER

Monday, 7 November 2022 16:10 (20 minutes)

Divertor detachment operation is of critical importance for future long pulse high power tokomaks, such as ITER and CFETR[1]. Partially detached divertor conditions are foreseen for ITER and DEMO. In conventional standard divertor (SD) operation, the effect of drifts on SD detachment is very strong, and the outer target heat loading is very higher than inner target with favorable-Bt discharge[2]. The root reason is the poloidal drift flux driving a lot of recycling plasma and radiated impurity into inner target region along private flux region. Currently, the experiment results have demonstrated that the drifts also have a great influence on the snowflake divertor detachment in TCV[3]. But, its detail physical mechanism is awaiting a better theoretical understanding of the relevant physical picture.

HL-2M allow operation with various advanced snowflake divertor configurations such as SF+ and SF-, and provides a good platform for studying advanced snowflake divertor plasma physics[4]. HL-2M snowflake divertor was designed by using SOLPS5.0[4]. The SOLPS5.0 modeling results without drifts showed that the inner target heat flux ($q_{\text{prep}} \sim 15 \text{ MWm}^{-2}$) and T_e ($T_e \sim 60 \text{ eV}$) are very higher than them in outer target plate ($q_{\text{perp}} \sim 1 \text{ MWm}^{-2}$, $T_e \sim 2 \text{ eV}$)[5]. Although the outer snowflake divertor with large magnetic expansion can effectively reduce the outer target heat loading, the inner divertor target heat loading is very high. In fact, the drifts may have a very stronger effect on snowflake divertor detachment/in-out asymmetry than SD. Therefore, in this work, we will investigate the influence of drifts on HL-2M snowflake divertor in-out target plate heat loading by employing the latest edge plasma code SOLPS-ITER[6] considering 100% all drifts and current.

References

- [1] A.S. Kukushkin, et al. Consequences of a reduction of the upstream power SOL width in ITER, J. Nucl. Mater. 438 (2013) 6–10.
- [2] V. Rozhansky, et al. Contribution of $E \times B$ drifts and parallel currents to divertor asymmetries, Nucl. Fusion. 52 (2012) 103017.
- [3] B.P. Duval, et al. Enhanced EXB drift effects in the TCV snowflake divertor, Nucl. Fusion. 55 (2015) 123023.
- [4] G.Y. Zheng, et al. Investigations on the heat flux and impurity for the HL-2M divertor, Nucl. Fusion. 56 (2016) 126013.
- [5] H. Du, et al. Exploring SF- in-out asymmetry and detachment bifurcation in HL-2M with $E \times B$ by SOLPS, Nucl. Mater. Energy. 22 (2020) 100719.
- [6] X. Bonnin, et al. ITER divertor plasma response to time-dependent impurity injection, Nucl. Mater. Energy. 0 (2016) 1–6.

Acknowledgments

This work was supported by National Natural Science Foundation of China under Grant Nos. 12175054; National Key R&D Program of China Nos. 2022YFE03180300, 2018YFE0301101.

Speaker's Affiliation

Southwestern Institute of Physics, ChengDu

Member State or IGO

China, People's Republic of

Primary author: DU, Hailong (Southwestern Institute of Physics)

Presenter: DU, Hailong (Southwestern Institute of Physics)

Session Classification: Poster Session I (DEMOs & Next Step Facilities)

Track Classification: Divertors for DEMO and Next-Step Facilities