



Contribution ID: 76

Type: Poster

## First results of the measurement of the $Ta(n,\gamma)$ cross-section at n\_TOF, CERN

Tuesday 9 July 2024 17:54 (1 minute)

V. Alcayne<sup>1</sup>, D. Cano-Ott<sup>1</sup>, E. González-Romero<sup>1</sup>, T. Martínez<sup>1</sup>, E. Mendoza<sup>1</sup>, A. Pérez de Rada Fiol<sup>1</sup>, A. Sánchez-Caballero<sup>1</sup>, J. Balibrea-Correa<sup>2</sup>, F. Calviño<sup>3</sup>, R. Capote<sup>4</sup>, A. Casanovas<sup>3</sup>, C. Domingo-Pardo<sup>2</sup>, J. LERENDUEGUI-MARCO<sup>2</sup> and the n\_TOF collaboration.

<sup>1</sup> Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Spain

<sup>2</sup> Instituto de Física Corpuscular, CSIC - Universidad de Valencia, Spain

<sup>3</sup> Universitat Politècnica de Catalunya, Spain

<sup>4</sup> International Atomic Energy Agency, Vienna-A-1400, PO Box 100, Austria

victor.alcayne@ciemat.es

Metallic alloys or metals with high melting points such as tantalum are being considered for the development of nuclear reactors for space. In recent critical experiments using highly enriched uranium or plutonium fuels, moderators and tantalum, large discrepancies have been found between the predicted and measured keff (i.e. needed critical masses). These observed discrepancies have been attributed to larger than reported uncertainties in the nuclear data of the materials involved, mainly tantalum, plutonium and graphite. The  $Ta(n,\gamma)$  cross section has also been reported as an important contributor to the uncertainty in the activation and heating of magnets used in large fusion reactors. The different measurements of the Ta neutron capture cross sections used in the evaluations are discrepant and affected by important experimental corrections like the self-shielding or angular correlations between  $\gamma$ -rays.

For these reasons, a new measurement of  $Ta(n,\gamma)$  cross section in the energy range from 0.1 eV to 500 keV has been performed at the n\_TOF facility. The use of C6D6 detectors at different angles and samples of different thicknesses have allowed us to overcome the limitations of the previous measurements. This abstract presents the experiment conducted at n\_TOF. It outlines the methodology employed for measurements and highlights the recent advancements in data analysis. In particular, the compatibility of the measurements made with different samples and different detectors will be presented as well as the preliminary comparison with the previous measurements and the most recent evaluations.

**Primary author:** ALCAYNE, Victor (CIEMAT)

**Co-authors:** SANCHEZ CABALLERO, Adrian (CIEMAT); PÉREZ DE RADA FIOL, Alberto (CIEMAT); Prof. CANO OTT, Daniel (CIEMAT - Centro de Investigaciones Energéticas, MedioAmbientales y Tecnológicas); MENDOZA CEMBRANOS, Emilio (CIEMAT); GONZÁLEZ ROMERO, Enrique (CIEMAT); CAPOTE NOY, Roberto Mario (IAEA NAPC-NDS); MARTINEZ PEREZ, Trino (CIEMAT); N\_TOF COLLABORATION

**Presenter:** SANCHEZ CABALLERO, Adrian (CIEMAT)

**Session Classification:** Poster Session

**Track Classification:** Measurements relevant to compound-nuclear reactions