

Recent advances about the unconformity-related U deposits

Tuesday, 24 June 2014 10:00 (30 minutes)

Giant unconformity-related uranium deposits were formed during the Mesoproterozoic era, 1.6-1.0 Ga ago, in both Athabasca Basin (Canada) and Kombolgie Basin (Australia). They are precious witnesses of protracted large-scale fluid flows at the interface between sedimentary basins and their crystalline basement, at conditions close to peak diagenesis (130-220°C). Although the Athabasca Basin hosts the world's largest high-grade uranium deposits, metallogenic models still bear important uncertainties. The objective of this contribution is to present new insights about the genetic model of these exceptional deposits.

Origin of the metals

The origin of the metals concentrated in the deposits has been investigated based on a systematic study of Hudsonian uranium mineralization in the basement rocks near the Athabasca Basin. This study shows that pre-Athabasca mineralization, expressed as uranium oxides, potentially represents a major uranium source for the unconformity related deposits.

Origin of the mineralizing brines

The origin of the brines has been investigated based on coupled Cl/Br and $\delta^{37}\text{Cl}$ composition of fluid inclusions trapped in quartz-dolomite veins and $\delta^{11}\text{B}$ composition of Mg-tourmalines associated with U ores. These studies have shown that the brines initially derive from subaerial evaporation of seawater up to epsomite saturation (salt content of ca. 25-35 wt%) forming the Cl-Na-K-Mg-rich brines.

Fluid percolation in the basement rocks, metal uptake and brine modifications

The original Cl-Na-K-Mg brines have percolated through the sedimentary pile and in the underlying basement, during tectonic reactivation, thanks to major faults and dense network of microfractures, partly inherited from late-orogenic deformation related to Trans-Hudson Orogen. Intensive brine/basement interaction was responsible for major chemical and isotopic changes (O, H, C) of the initial brines to form two chemically distinct NaCl-rich and CaCl₂-rich brines, both being highly enriched in metals. Their metal enrichment comparable with brines related to MVT Pb-Zn deposits supports the idea that the basement was the dominant source for metals, and especially for U.

Conditions for the transport and deposition of uranium

The mineralizing brines have U concentrations between 1×10^{-6} and 2.8×10^{-3} mol.l⁻¹, making them the U richest crustal fluids so far. This exceptional U content is related to the oxidizing and acidic nature of the brines and to the high availability of U sources. Synchrotron analyses show that U is present at the hexavalent state in these brines. The mixing of the NaCl-rich and CaCl₂-rich brines is coeval with the UO₂ deposition but the reductant necessary for UO₂ precipitation remains enigmatic.

Primary author: Dr MERCADIER, Julien (Georessources)

Co-authors: Dr RICHARD, Antonin (GeoRessources); Dr ANNESLEY, Irvine R. (Department of Geological Sciences, University of Saskatchewan); Dr BOIRON, Marie-Chirstine (GeoRessources); Dr CATHELINÉAU, Michel (GeoRessources); Dr CUNÉY, Michel (CNRS - GeoRessources - CREGU - Université de Lorraine)

Presenter: Dr MERCADIER, Julien (Georessources)

Session Classification: Uranium geology

Track Classification: Uranium geology