

### Updates from OCL

CM on Evaluation of Photon Strength Function



Funded by The Research Council of Norway

V. W. Ingeberg

### Outline


- Recently published and soon to be published
- Ongoing work
- Recent experiments




# OM measurements





The Oslo Method experiment






### Oslo method experiment





### Oslo Method in 4 (simple) steps!

- 1. Obtain excitation energy tagged gamma spectra
- 2. Unfolding method
- 3. First generation method
- 4. Extract NLD and  $\gamma$ SF



$$P(E_x, E_\gamma) \propto 2\pi E_\gamma^3 f(E_\gamma) \rho(E_f = E_x - E_\gamma)$$



UNIVERSITY

## Unfolding

1. 
$$f^0 = \mathbf{R} u^0$$

2. 
$$u^1 = u^0 + (m - f^0)$$


Repeat until convergence

f: Folded spectrum

u: unfolded

m: measured spectrum


**R**: response function




**UNIVERSITY** 

### Sources of errors – unfolding method

Assuming a efficiency reduction of ≈ 20%









### Uncertainty estimation

### Regularized Unfolding of $\gamma$ -ray Spectra for Nuclear Physics Applications

E. Lima, 1, 2 L. L. Braseth, 1 A. H. Mjøs, 1, 2 M. Hjorth-Jensen, 3, 4 A. Kvellestad, 1 and A. C. Larsen 1, 2 <sup>1</sup>Department of Physics, University of Oslo, N-0316 Oslo, Norway <sup>2</sup>Norwegian Nuclear Research Centre, Norway\* <sup>3</sup> Department of Physics and Center for Computing in Science Education, University of Oslo, N-0316 Oslo, Norway <sup>4</sup>Department of Physics and Astronomy and Facility for Rare Isotope Beams, Michigan State University, East Lansing, MI 48824, USA (Dated: June 5, 2025)







### Bayesian Unfolding in the Oslo Method

### Model outline:


- Factorize total detector response R: discrete response D and Gaussian response kernel G from the finite detector resolution
- Forward model:  $\mu \to \eta = G\mu \to \nu = D\eta$ ( $\mu$  = counts in true space,  $\eta$  = counts smeared by G,  $\nu$  = expected counts in observed space)
- Likelihood:  $n_i \sim Poisson(v_i)$  captures counting-statistics uncertainty
- Prior: log-normal on  $\mu$  with hierarchical hyperpriors
- Posterior exploration with No-U-Turn Sampler (NUTS, PyMC 5)
- Capabilities & benefits
  - Supports spectra up to 1000 energy bins
  - Delivers full Bayesian uncertainty bands per bin (+ correlation between bins)
  - Reproducible: open-source code & data
- Planned journal submission: early autumn 2025
- Co-authors: Erlend Lima, Ann-Cecilie Larsen, Anders Kvellerstad, Morten Hjort-Jensen



### Extraction of NLD and GSF

Higher order estimates through a least  $\chi^2$ -minimization:

$$P_{th}(E_x, E_{\gamma}) = \frac{2\pi E_{\gamma}^3 f(E_{\gamma}) \rho(E_x - E_{\gamma})}{\sum_{E_{\gamma}} 2\pi E_{\gamma}^3 f(E_{\gamma}) \rho(E_x - E_{\gamma})}$$



$$\chi^{2} = \frac{1}{N_{free}} \sum_{E_{i}=E_{min}}^{E_{max}} \sum_{E_{\gamma}=E_{\gamma}^{min}}^{E_{i}} \left[ \frac{P_{th}(E_{i},E_{\gamma}) - P(E_{i},E_{\gamma})}{\Delta P(E_{i},E_{\gamma})} \right]^{2}$$

Each vector element in  $\rho$  and f is treated as a free parameter

56Fe(p,p') example:

Data points ("pixels"): 2052

Free parameters: 184

 $N_{free} \ll N_{data}$ 

Runs 50 iterations, but converges often at 10-20 iterations (depending on the case)


24.11.2025

### Normalization of NLD

- Comparison to NLD from tabulated discrete levels
- Comparison to NLD at neutron separation energy

$$\rho(S_n) = \frac{\sigma^2}{D_0} \frac{2}{\alpha_{\pi} + 1} \frac{1}{(I_t + 1) \exp(-(I_t + 1)^2 / 2\sigma^2) + I_t \exp(-I_t^2 / 2\sigma^2)}$$

- $D_0$ : Average s-wave resonance spacing
- $\alpha_{\pi}$ : Parity asymmetry
- σ: Spin-cutoff parameter
- Between S<sub>n</sub> and data: Interpolation with some model (usually CT)
- Gives the absolute value of the NLD, A, and the slope  $\alpha$





### Normalization of NLD – NLD at particle threshold

Use resonance spacing at particle threshold to determine NLD:

$$\frac{1}{D_0} = \frac{1}{2} \left[ \rho(S_n, I_t + 1/2) + \rho(S_n, I_t - 1/2) \right]$$

$$\rho(S_n) = \frac{\sigma^2}{D_0} \frac{2}{\alpha_\pi + 1} \frac{1}{(I_t + 1) \exp(-(I_t + 1)^2/2\sigma^2) + I_t \exp(-I_t^2/2\sigma^2)}$$

- Where:
  - $D_0$ : s-wave resonance spacing, most often neutron resonaces, sources such as RIPL-3 and Mughabghab
  - *I<sub>t</sub>*: Ground state spin of the A-1 nucleus
  - $\alpha_{\pi}$ : Parity asymmetry
  - $\sigma$ : Spin-cutoff parameter



### Normalization of GSF

$$\tilde{\rho}(E_i - E_{\gamma}) = A\rho(E_i - E_{\gamma})e^{\alpha(E_i - E_{\gamma})}$$
$$\tilde{\mathcal{T}}(E_{\gamma}) = B\mathcal{T}(E_{\gamma})e^{\alpha E_{\gamma}}$$

Slope parameter  $\alpha$  determined by normalized NLD

Now assuming dipole transitions dominate:

$$BT(E_{\gamma}) = B \sum_{XL} \mathcal{T}_{XL}(E_{\gamma}) \approx B[\mathcal{T}_{E1}(E_{\gamma}) + \mathcal{T}_{M1}(E_{\gamma})]$$

Next step: Average total radiative width:

$$\langle \Gamma_{\gamma}(E,J,\pi) \rangle = \frac{1}{4\pi\rho(E,J,\pi)} \sum_{XL} \sum_{I_f,\pi_f} \int_0^E dE_{\gamma} \, \mathcal{T}_{XL}(E_{\gamma}) \rho(E-E_{\gamma},J_f,\pi_f)$$

Normalization integral (normalize to  $\langle \Gamma_{\gamma 0} \rangle$ , average radiative of s-wave capture res.)

$$\langle \Gamma_{\gamma} \left( S_n, I_t \pm \frac{1}{2}, \pi_t \right) = \frac{BD_0}{2\pi} \int_0^{S_n} dE_{\gamma} \, \mathcal{T}_{XL} (E_{\gamma}) \rho \left( S_n - E_{\gamma} \right) \, \sum_{J=-1}^1 g(S_n - E_{\gamma}, I_t \pm \frac{1}{2} + J)$$




### Normalization of GSF


- GSF covers energies between  $E_{\nu}^{min}$  and  $E_{x}^{max}$
- Need to extrapolate between
  - $E_{\gamma} = 0$  and  $E_{\gamma}^{min}$
  - $E_x^{max}$  and  $S_n$
- Between  $E_{\gamma} = 0$  and  $E_{\gamma}^{min}$ :

$$\mathcal{T}(E_{\gamma}) = \frac{\exp(aE_{\gamma} + b)}{E_{\gamma}^{3}}$$

• Between  $E_x^{max}$  and  $S_n$ :

$$\mathcal{T}(E_{\gamma}) = \exp(cE_{\gamma} + d)$$







### Systematic effects of the normalization

- Selection of spin-cutoff parameter:
  - If spin-cutoff is to big -> steep slope
  - Will affect the strength abs. value
- Large parity asymmetry
  - Will give wrong abs. Strength
  - Average total radiative width of s-wave resonances integral no longer valid

$$\langle \Gamma_{\gamma}(S_n, I_t \pm 1/2, \pi_t) \rangle = \frac{B}{2\pi \rho(S_n, I_t \pm 1/2, \pi_t)}$$

$$\times \int_0^{S_n} dE_{\gamma} \left[ \mathcal{T}_{E1}(E_{\gamma}) \mathcal{P}_{\mp} + \mathcal{T}_{M1}(E_{\gamma}) \mathcal{P}_{\pm} \right]$$


$$\times \rho(S_n - E_{\gamma}) \sum_{I=1}^1 g(S_n - E_{\gamma}, I_t \pm 1/2 + J)$$



### Shape method

- Decay to discrete levels
- Can obtain the shape of the  $\gamma$ SF


• Constrains the slope of OM data 
$$N_D \propto f(E_\gamma) E_\gamma^3 \sum_{[J_f]}^{J_i = J_f + 1} \sigma(E_i, J_i) g(E_i, J_i).$$
 
$$f(E_{\gamma 1}) \propto \frac{N_{D1}}{E_\gamma^3 \sum_{[J_f 1]} \sum_{J_i = J_f - 1}^{J_i = J_f + 1} g(E_i, J_i)}$$
 (a)  ${}^{92}Zr(p, p'\gamma)^{92}Zr$   $D_4$   $D_3$   $D_2$   $D_1$   $10^3$  
$$f(E_{\gamma 2}) \propto \frac{N_{D2}}{E_\gamma^3 \sum_{[J_f 2]} \sum_{J_i = J_f 2 - 1}^{J_i = J_f 2 + 1} g(E_i, J_i)}$$
 
$$g(E_i, J_i) = \frac{N_{D2}}{N_D}$$
 
$$g(E_i, J_i) = \frac{N_{D3}}{N_D}$$
 
$$g(E_i, J_i) = \frac{N_D}{N_D}$$
 
$$g(E_i,$$






### Reaction dependent systematic errors

- (d, p) reaction
  - Close to Coulomb barrier – reaction resembles a deuteron breakup + neutron absorption
  - Limited spin window populated
- Test with RAINIER (DICEBOX)



### Theoretical spin-parity population Green's function transfer formalism






UNIVERSITY

Reaction dependent systematic errors

### Nd chain: (p,p) and (d,p)


- Limited spin population
- Shape method indicates that OM overestimates the slope significantly
- For (d,p): 11% of full NLD observed
- For (p,p): 22% of full NLD observed
- Independently confirmed by measurements of ground state band feeding
- M. Guttormsen et al., Phys. Rev. C 106, 034314 (2022)






### Reaction Spesific Errors


 Initial population determined by side-feeding








### Recent results




V. W. Ingeberg *et al.*, Phys. Rev. C **111**, 015803 (2025)



A. C. Larsen et al., Phys. Rev C 108, 025804 (2023)



UNIVERSITY



<sup>124</sup>Sn

16 18

10

 $E_{\gamma}$  (MeV)

12

120Sn

16

14

8

6

10

E<sub>γ</sub> (MeV)

18 0

10



### Recent experiments at OCL

93Nb + 16-MeV protons, April 2025 a)

PI: Sivahami Uthayakumaar (FRIB), Ann-Cecile Larsen

**Motivation:** The  $^{92}$ Nb(n,y) $^{93}$ Nb is one of the most important reactions for the destruction of  $^{92}$ Nb in the p-process nucleosynthesis. We aim to constrain this un-measured reaction cross section and astrophysical reaction rate by using the level density and gamma strength function from the Oslo method on the  $^{93}$ Nb(p,p' $^{\gamma}$ ) data set.

<sup>106</sup>Pd, <sup>59</sup>Co, <sup>27</sup>Al + 16-MeV protons, June 2025 b)

PI: Vetle W. Ingeberg, Ann-Cecilie Larsen, Kevin Li

**Motivation:** The PSF of <sup>59</sup>Co was recently published from a NRF experiment and has an unusual shape. The <sup>27</sup>Al is of interest in supernova nucleosynthesis and are close to the limit of the Oslo method. In <sup>106</sup>Pd there are discrepancies in the nuclear level density and between the models of the NLD.

<sup>154</sup>Sm + 30 MeV alphas. August 2025 c)

PI: Lauren T. Bell. Sunniva. Ann-Cecilie Larsen

Motivation: We want to to investigate the impact of angular-momentum transfer on the level density and gamma strength function of <sup>154</sup>Sm by comparing the <sup>154</sup>Sm(a,a'q) data to previously taken data on <sup>154</sup>Sm(p,p'q) Bonus: We also got data on the exotic, n-rich 155,156Eu through the <sup>154</sup>Sm(a,tg)<sup>155</sup>Eu and <sup>154</sup>Sm(a,dg)<sup>156</sup>Eu reaction channels, respectively, of interest for the intermediate neutron-capture process

<sup>130</sup>Te + 30 MeV alphas & <sup>130</sup>Te + 24 MeV alphas, September 2025 d)

PI: Claudia, Therese (Master students), Ann-Cecilie, Maria

Motivation: to get level densities and gamma strength for <sup>132</sup>I and <sup>131</sup>I through the reaction channels <sup>130</sup>Te(a,pq)<sup>133</sup>I, <sup>130</sup>Te(a,dq)<sup>132</sup>I, <sup>130</sup>Te(a,tq)<sup>131</sup>I respectively, for the i-process nucleosynthesis Bonus: we also have a super nice 130Te(a,a'g) data set, for nuclear structure (oblate deformation, looking for the M1 scissors mode in an oblate nucleus for the first time)

<sup>116</sup>Cd + 30 MeV alphas. October 2025 e)

PI: Maria Markova. Ann-Cecile Larsen

Motivation: to get level densities and gamma strength for 116Cd through the reaction channel 116Cd(a,a'g)116Cd, to get a data-constrained 115Cd(n,g) reaction rate, of critical importance for the i-process nucleosynthesis Bonus: we also got data on the exotic, n-rich 117,118 In through the 116Cd(a,tg)117 In and 116Cd(a,dg)118 In reaction channels, respectively, also of interest for the intermediate neutron-capture process, and to investigate the pygmy dipole resonance for indium isotopes, as 109 In shows almost no sign of a PDR

<sup>107,109</sup>Aa + 16-MeV protons. November 2025

PI: Henrik D. Andrews, Ann-Cecilie Larsen, Sunniva Siem

Motivation: Study level densities and gamma strength functions in silver isotopes for the first time. Search for the pygmy dipole resonance. Also, i-process relevant


<sup>186</sup>W + 30 MeV alphas. November 2025 g)

PI: Rahul Jain (LLNL), Richard Hughes (LLNL), Andrea Richard (Ohio Univ.), Ann-Cecilie Larsen


Motivation: Aim was to reach 189Re and 188Re with the 186W(a.p) and 186(a.d) reactions.



### GSF of <sup>59</sup>Co



Will measure with both (p,p') and (a,a')



~ 2-3 hours of data

# SUPER Preliminary


R. Schwengner et al., Phys. Rev. C 111, 014312 (2025)




UNIVERSITY

# Limits of the Oslo Method • MSc project of Lise Granheim • Can you extract GSF from 280:

- <sup>28</sup>Si
- Can you extract GSF and NLD in <sup>27</sup>Al?
  - We have measured (p, p')






 $E_{\gamma}$  (MeV)




UNIVERSITY OF OSLO

### New project on Ag



Plan to investigate Ag nuclei with several probes

PhD project of Henrik Andrews





# AVIT application, StarLight prototype

- 3 hexagonal-shaped scintillator crystals made of NaI(TI+Li) that can separate neutrons and gammas through pulse-shape discrimination
- Testing &characterization at OCL
- Full array financed hopefully through a grant in 2026/2027 [PI Ann-Cecilie Larsen]

