



IAEA FEC 2014

Contribution ID: 106

Type: **Poster**

Computational Study of Magnetic Field Compression by Laser Driven Implosion

Thursday, 16 October 2014 14:00 (4h 45m)

A compression of external magnetic field by a laser driven implosion is studied using two dimensional radiation hydrodynamic simulation. The simulation results show that (i) it is possible to compress the magnetic field to be 10kT (108 Gauss), and (ii) the strong magnetic field should affect the implosion dynamics because of the suppression of the electron heat flux which across strong magnetic field lines. This result suggest that a target and initial conditions for fast ignition with external magnetic field must be designed carefully not only for the control of the hot electron transport but also for the formation of high dense plasma.

Paper Number

IFE/P6-6

Country or International Organisation

Japan

Primary author: Mr NAGATOMO, Hideo (Osaka University)

Co-authors: Prof. NISHIGUCHI, Akio (Osaka Institute of Technology); Dr SUNAHARA, Atsushi (Institute for Laser Technology); Prof. AZECHI, Hiroshi (Institute of Laser Engineering, Osaka University); Prof. SHIRAGA, Hiroyuki (Institute of Laser Engineering, Osaka University); Prof. SAKAGAMI, Hitoshi (National Institute for Fusion Science); Prof. MIMA, Kunioki (Graduate School for the Creation of New Photonics Industries); Dr FUJIOKA, Shinsuke (Institute of Laser Engineering, Osaka University); Dr JOHZAKI, TOMOYUKI (Graduate School of Engineering, Hiroshima University); Dr SANO, Takayoshi (Institute of Laser Engineering, Osaka University)

Presenter: Mr NAGATOMO, Hideo (Osaka University)

Session Classification: Poster 6