



IAEA FEC 2014

Contribution ID: 144

Type: Poster

## Self Organization of High $\beta_p$ Plasma Equilibrium with an Inboard Poloidal Null Sustained by Fully Non-Inductive Current Drive in QUEST

*Tuesday 14 October 2014 08:30 (4 hours)*

There is a considerable interest in operating tokamak at high value of  $\beta_p$ , but maximum attainable  $\beta_p$  is limited by a equilibrium limit with appearance of a null point at the inboard side. Such configuration is realized transiently earlier by Electron Cyclotron Waves, but in QUEST such equilibrium is stably produced in steady state and its equilibrium properties are investigated. In QUEST, successful production of high  $\beta_p$  plasma ( $\beta_p > 1$ ) and its long pulse sustainment by fully non-inductive (NI) current drive with the help of a modest power ( $< 100$  kW) ECW is demonstrated. High  $\beta_p$  plasma is formed by confining energetic electrons produced by multiple resonant EC interaction in a high magnetic mirror configuration and high  $B_z/B_t > 0.1$ . We found that (i) high  $\beta_p$  plasma is naturally self organized to form a stable natural Inboard Poloidal field Null (IPN) equilibrium, (ii) a critical  $\beta_p$ , which defines the transition boundary from Inboard Limiter (IL) to IPN equilibrium and (iii) a new feature of plasma self organization to enhance its negative triangular shape to sustain high  $\beta_p$ . With high  $\beta_p$  formation, plasma naturally self organizes itself to reduce the elongation as first observed in TFTR, which is also observed in the present case. However, we found a new self organization feature, where plasma shape adjusts itself to become more negatively triangular. This new feature overcompensates the diminution of  $\beta_p$  due to the reduction in elongation. A simple analytic solution of Grad-Shafranov equation is applied to investigate such aspect. The model supports this facet which, predicts higher  $\beta_p$  at larger negative triangularity. The boundary flux surfaces generated through the model agree well with the measurements. The model is also in agreement with the critical  $\beta_p$  for IL to IPN transition, which is very well matched with the measurements. This result shows a relatively simple method to produce and sustain high  $\beta_p$  plasma close to the equilibrium limit in a stable configuration exploiting its self organization property.

### Country or International Organisation

Japan

### Paper Number

EX/P1-39

**Author:** Mr MISHRA, kishore (Kyushu University)**Co-authors:** ZUSHI, Hideki (Riam Kyushu University); Dr IDEI, Hiroshi (Research Institute for Applied Mechanics, Kyushu University); Dr HASEGAWA, Makoto (RIAM, Kyushu University); Dr ONCHI, Takumi (RIAM, Kyushu University)**Presenter:** Mr MISHRA, kishore (Kyushu University)

**Session Classification:** Poster 1