



IAEA FEC 2014

Contribution ID: 240

Type: Poster

## Development of Divertor Simulation Research in the GAMMA 10/PDX Tandem Mirror

Friday 17 October 2014 14:00 (4h 45m)

This paper presents the first achievement of detachment from high temperature plasma equivalent to the SOL plasma of tokamaks using a large tandem mirror device. In Plasma Research Center, University of Tsukuba, we have started divertor simulation experiments at the end-cell of GAMMA 10/PDX (E-divertor) in order to realize the divertor simulation closely resemble to actual divertor environment [1, 2].

The first experiment for realizing detached plasma state from the high-temperature plasmas has been performed using H<sub>2</sub> and noble gas injection in the divertor simulation experimental module (D-module) recently installed in the west end-cell of GAMMA 10/PDX. Here, the plasma with  $n_e \sim 2 \times 10^{18} \text{ m}^{-3}$  and  $T_{i//} \sim 150 \text{ eV}$  was produced with two ICRF waves (RF1 and RF2) in the upstream region (central-cell). The highest electron density of  $2.4 \times 10^{18} \text{ m}^{-3}$  was obtained by simultaneous injection of H<sub>2</sub> and Xe into D-module. The particle and heat fluxes on the V-shaped target plate mounted in D-module were measured and the dependence on the Ar plenum pressure was examined. As increasing the amount of injection gas, both particle and heat fluxes continuously decrease. It is also observed that the electron temperature is drastically reduced from few tens eV to  $\sim 3 \text{ eV}$  due to the Ar injection. Furthermore, in the case with H<sub>2</sub> and Xe, the ion flux was almost fully suppressed at the V-shaped corner. At the same time, 2-D image of H $\alpha$  emission captured with the fast camera showed the clear detachment of emission zone from the target plate. Above results clearly show the evidence of plasma detachment in D-module.

Recently two antennas for additional ICRF heating (RF3) were installed in anchor cells in order to build up both particle and heat fluxes at the end-cell. A remarkable increase of the plasma density and end-loss ion flux is achieved during RF3. We evaluated the particle flux at the mirror exit by investigating correlation between particle flux determined from the probe measurements near the mirror throat and the end-loss ion current measured with end-loss ion energy analyzer (ELIEA). From this evaluation, the particle flux is estimated to be  $1.7 \times 10^{23} / \text{m}^2 \text{ s}$  at the end-mirror exit.

[1] Y. Nakashima, et al., Fusion Eng. Design volume 85 issue 6 (2010) 956-962.

[2] Y. Nakashima, et al., Journal of Nuclear Materials 438 (2013) S738-S741.

### Country or International Organisation

Japan

### Paper Number

FIP/P8-10

**Primary author:** Prof. NAKASHIMA, Yousuke (Plasma Research Center, University of Tsukuba)

**Co-authors:** Prof. SAGARA, Akio (National Institute for Fusion Science); Prof. TONEGAWA, Akira (Tokai University); Prof. HATAYAMA, Akiyoshi (Keio University); Dr KUBO, Hiroataka (Japan Atomic Energy Agency); Dr MATSUURA, Hiroto (Osaka Prefecture University); Mr TAKEDA, Hisato (Plasma Research Center, University of Tsukuba); Prof. KATANUMA, Isao (University of Tsukuba); Dr KOHAGURA, Junko (Plasma Research Center,

University of Tsukuba); Mr HOSOI, Katsuhiro (Plasma Research Center, University of Tsukuba); Mr ICHIMURA, Kazuya (Plasma Research Center, University of Tsukuba); Dr OKI, Kensuke (Plasma Research Center, University of Tsukuba); Dr HIRATA, Mafumi (Plasma Research Center, University of Tsukuba); Prof. ICHIMURA, Makoto (Plasma Research Center, University of Tsukuba); Dr SHOJI, Mamoru (National Institute for Fusion Science); Dr FUKUMOTO, Masakatsu (Japan Atomic Energy Agency); Dr YOSHIKAWA, Masayuki (University of Tsukuba); Ms IWAMOTO, Miki (Plasma Research Center, University of Tsukuba); Prof. SAKAMOTO, Mizuki (Plasma Research Center, University of Tsukuba); Dr NISHINO, Nobuhiro (Hiroshima University); Mr ASAKURA, Nobuyuki (Japan Atomic Energy Agency); Prof. OHNO, Noriyasu (Nagoya University); Dr MINAMI, Ryutaro (Plasma Research Center, University of Tsukuba); Dr IKEZOE, Ryuya (Plasma Research Center, University of Tsukuba); Dr KADO, Shinichiro (Kyoto University); Dr NAGATA, Shinji (Tohoku University); Dr MASUZAKI, Suguru (National Institute for Fusion Science); Prof. SHIKAMA, Tatsuo (Tohoku University); Dr NUMAKURA, Tomoharu (Plasma Research Center, University of Tsukuba); Prof. IMAI, Tsuyoshi (Plasma Research Center, University of Tsukuba); Dr KARIYA, Tsuyoshi (Plasma Research Center, University of Tsukuba); Mr HOSODA, Yasunari (Plasma Research Center, University of Tsukuba); Prof. HIROOKA, Yoshihiko (National Institute for Fusion Science); Prof. UEDA, Yoshio (Osaka University)

**Presenter:** Prof. NAKASHIMA, Yousuke (Plasma Research Center, University of Tsukuba)

**Session Classification:** Poster 8