



IAEA FEC 2014

Contribution ID: 162

Type: Poster

Fast Ignition Experiments and Intense Hard-X-Ray Harsh Environment

Thursday, 16 October 2014 14:00 (4h 45m)

Fast ignition (FI) experiments have been performed at the Institute of Laser Engineering, Osaka University by using Gekko-XII laser for implosion and LFEX laser for fast heating of fuel targets. In FI experiment with intense heating laser irradiation ($>10^{19}$ W/cm²), most of heating laser energy is converted to that of hot electrons with energy ranging well up to several 10 MeV. Intense hard x-rays (or so-called “ γ rays” in terms of the photon energy) are generated from hot electrons via Bremsstrahlung in the target material. Such high-energy photons are a real threat as intense background signals to the x-ray and neutron diagnostics. Neutrons generated via (γ ,n) reactions taking place in and around the target chamber are also a threat to the neutron diagnostics. We found that FI experiments are in such a γ -ray and neutron harsh environment, and evaluated those conditions which have never been experienced in conventional laser-plasma experiments. We have developed many new plasma diagnostics that are compatible with such a γ -ray and neutron harsh environment. FI physics- and integrated-experiments have been successfully performed with those improved plasma diagnostics with much better accuracy than in the previous experiments even in such a γ -ray and neutron harsh environment. These works are also useful as test pilot experiments for future experimental fusion reactors in which much more harsh environment is anticipated.

Paper Number

IFE/P6-3

Country or International Organisation

Japan

Primary author: Prof. SHIRAGA, Hiroyuki (Institute of Laser Engineering, Osaka University)

Co-authors: Prof. NISHIMURA, Hiroaki (Institute of Laser Engineering, Osaka University); Prof. AZECHI, Hiroshi (Institute of Laser Engineering, Osaka University); Prof. MAYUKO, Koga (University of Hyogo); Prof. NAKAI, Mitsuo (Institute of Laser Engineering, Osaka University); Prof. FUJIOKA, Shinsuke (Institute of Laser Engineering, Osaka University); Prof. OZAKI, Tetsuo (National Institute for Fusion Science); Dr ARIKAWA, Yasunobu (Institute of Laser Engineering Osaka University)

Presenter: Prof. SHIRAGA, Hiroyuki (Institute of Laser Engineering, Osaka University)

Session Classification: Poster 6