



IAEA FEC 2014

Contribution ID: 679

Type: Poster

## Modelling of Transitions Between L- and H-Mode Including W Behaviour in ITER Scenarios

Wednesday 15 October 2014 08:30 (4 hours)

The dynamics of the access to and exit from high QDT regimes in the H-mode confinement regime in ITER is expected to be qualitatively different to present experiments: neutral fuelling is much less effective,  $P_{sep}/PL-H < 2.0$  even in stationary QDT  $\sim 10$  burning conditions, the density evolution determines not only PL-H but also  $P_{alpha}$  which in turn affects  $dW_{th}/dt$  after a transition, and plasma position control may be challenging in case of an unexpected back transition to L-mode. In addition, the presence of W may impose additional operational constraints due to possible core accumulation and increased radiation during transients (possibility of a sudden return to L-mode confinement, plasma-wall contact and/or a disruption). To determine under which conditions the transition to stationary high QDT H-mode regime and its safe termination can be achieved, how the plasma evolution to/from H-mode can be optimised, and to assess the problem of possible core W accumulation, modelling studies have been carried out with the JINTRAC suite of codes, simulating the core and core+SOL plasma evolution for the entire period of density evolution following transitions to/from H-mode in the ITER 15 MA/5.3 T and 7.5 MA/2.65 T scenarios.

Simulation scans for the L-H transition have been performed with varying target waveforms for the density evolution, applying a feedback on pellet fuelling. Depending on boundary and operational conditions, limits for the density ramp rate and/or a delay time before the application of increased fuelling could be established. Below these limits, the plasma remains in dithering conditions with  $P_{sep}-PL-H$  for a long while before it enters a good quality H-mode regime at  $P_{sep}>PL-H$ , leading to increased flux consumption and a significantly reduced burn duration. In extreme cases, the plasma never reaches high performance H-mode and returns back to L-mode.

The back transition to L-mode has also been assessed. The fast reduction in core energy would cause  $P_{sep}$  to remain close to PL-H. The plasma would then not immediately reach L-mode but stay in H-mode for a while, followed by a dithering phase before the ETB completely disappears. The energy loss could become accelerated though by an immediate transition to dithering mode e.g. after a strong MHD event. Subsequent W accumulation could then lead to an immediate transition to L-mode and a disruption.

### Country or International Organisation

Austria

### Paper Number

TH/P3-24

**Author:** Dr KOECHL, Florian (Vienna University of Technology, Institute of Atomic and Subatomic Physics)

**Co-authors:** Dr LOARTE, Alberto (ITER Organization); Dr POLEVOI, Alexei (ITER Organization); Dr KUKUSHKIN,

Andrei S. (ITER Organization); Dr SAIBENE, Gabriella (Fusion for Energy Joint Undertaking (F4E)); Dr SARTORI, Roberta (Fusion for Energy Joint Undertaking (F4E)); Dr PARAIL, Vassili (CCFE)

**Presenter:** Dr KOECHL, Florian (Vienna University of Technology, Institute of Atomic and Subatomic Physics)

**Session Classification:** Poster 3