

Multi-Scale ITG/TEM/ETG Turbulence Simulations with Real Mass Ratio and β Value

S. Maeyama, Y. Idomura, M. Nakata,
M. Yagi, N. Miyato
Japan Atomic Energy Agency

Collaborators: T.-H. Watanabe (Nagoya Univ.),
M. Nunami, A. Ishizawa (NIFS)



25th IAEA FEC, 15 Oct. 2014




Introduction

One of the critical issues in ITER is electron heat transport, which is inherently multi-scale physics.

2000 [Jenko00PoP]

A candidate is electron temperature gradient modes (**ETG**).

“Streamers” in ETG turbulence



2007 [Candy07PPCF, Waltz08PoP, Görler08PRL]

ETGs give small transport if there are ion temperature gradient and trapped electron modes (**ITG/TEM**).

However, these multi-scale simulations were limited:

- Reduced mass ratio ($m_i/m_e = 400, 900$)
- Electrostatic approximation ($\beta = 0$)

Motivation & Outline


Following points are not yet clarified:

- (i) Are there multi-scale interactions even **with the real mass ratio** and β value?

- (ii) If yes, how do the interactions occur?

Linear instabilities from electron to ion scales

Scale separation with real mass ratio




Motivation & Outline

Following points are not yet clarified:

- (i) Are there multi-scale interactions even **with** the real mass ratio and β value?
 - **Multi-scale simulation demonstrates cross-scale interactions.**

- (ii) If yes, how do the interactions occur?
 - **Nonlinear interaction analysis reveals their mechanisms.**

Linear instabilities from electron to ion scales




The GKV code

[Watanabe06NF,Maeyama13CPC]

- Solve gyrokinetic ions and electrons with electromagnetic fluctuations in a flux-tube geometry.
- Validation with experiments. [Posters:Nakata,Ishizawa,Nunami]
- High scalability allows ITG/TEM/ETG simulations with ~100k CPU cores in ~100 hours. [Maeyama13SC]


Plasma parameters are
Cyclone base case
parameters [Dimits00PoP]

- $R/L_{Ti}=R/L_{Te}=6.82$,
 $R/L_n=2.2$, $T_e=T_i$,
 $r/R=0.18$, $q=1.4$,
 $s=0.786$
- Real mass ratio:
 $m_i/m_e=1836$
- Real β value: $\beta=2.0\%$
(below NZT [Pueschel13PRL])





Multi-scale turbulence simulation ($\beta=2.0\%$)

Time evolution of the electrostatic potential fluctuations
(at mid-plane of the flux tube)





Energy spectrum

($\beta = 2.0\%$)






Energy spectrum

($\beta = 2.0\%$)





Energy spectrum

($\beta = 2.0\%$)





Energy spectrum

($\beta = 2.0\%$)





Energy spectrum

($\beta = 2.0\%$)





Energy spectrum

($\beta = 2.0\%$)



Electron energy diffusion spectrum in multi-scale turbulence is NOT a sum of single-scale ones.

- In zero- β case, due to strong electron-scale suppression, ion-scale simulations give a good estimate.
- In finite- β case, electron-scale suppression is weak. Ion-scale transport is enhanced in multi-scale analysis.



* Ion energy diffusion is similar (see proceedings).

Analysis of nonlinear interactions

Gyrokinetic triad transfer


- Mode-to-mode nonlinear transfer of perturbed entropy

$$I_{sk} = \sum_p \sum_q J_{sk}^{p,q} \quad [\text{Nakata12PoP}]$$

$$J_{sk}^{p,q} = \delta_{k+p+q,0} \frac{\mathbf{b} \cdot \mathbf{p} \times \mathbf{q}}{2B} \times \text{Re} \left[\left\langle \int d\nu^3 (\bar{\chi}_{sp} g_{sq} - \bar{\chi}_{sq} g_{sp}) \frac{T_s g_{sk}}{F_{SM}} \right\rangle \right]$$

(Generalized potential $\bar{\chi}_{sk} = (\bar{\phi}_k - v_{\parallel} \bar{A}_{\parallel k})$,

Nonadiabatic distribution $g_{sk} = f_{sk} + \frac{e_s F_{SM}}{T_s} \bar{\phi}_k$)



Electron-scale suppression mechanism:

- **Ion-scale ZF shearing? Or, Another structures?**


Ion-scale enhancement mechanism:

- **Inverse cascade to ion-scale turbulence?
Or, Damping of ion-scale ZFs?**

Suppression of electron-scale streamers by high- k_x ITG/TEM structures.


Triad transfer $\sum_s J_{sk}^{p,q}$ for a streamer

$(k_x p_{ti}, k_y p_{ti}) = (0, 4.4)$ at $t = 20 - 30 R/v_{ti}$




Suppression of electron-scale streamers by high- k_x ITG/TEM structures.

Triad transfer $\sum_s J_{sk}^{p,q}$ for a streamer
 $(k_x p_{ti}, k_y p_{ti}) = (0, 4.4)$ at $t = 20 - 30R/v_{ti}$




Kinetic electrons create fine radial structures ($k_x p_{ti} > 1$). [Dominski12JPCS, Maeyama14PoP]




Suppression of electron-scale streamers by high- k_x ITG/TEM structures.

Triad transfer $\sum_s J_{sk}^{p,q}$ for a streamer
 $(k_x p_{ti}, k_y p_{ti}) = (0, 4.4)$ at $t = 20-30R/v_{ti}$




Kinetic electrons create fine radial structures ($k_x p_{ti} > 1$). [Dominski12JPCS, Maeyama14PoP]

At the reduction phase, these radial structures suppress streamers.




Suppression of electron-scale streamers by high- k_x ITG/TEM structures.

Triad transfer $\sum_s J_{sk}^{p,q}$ for a streamer
 $(k_x p_{ti}, k_y p_{ti}) = (0, 4.4)$ at $t = 20-30R/v_{ti}$




Kinetic electrons create fine radial structures ($k_x p_{ti} > 1$). [Dominski12JPCS, Maeyama14PoP]

At the reduction phase, these radial structures suppress streamers.




After suppression of ETG/streamers, normal cascade dominates electron scales.

Triad transfer $\sum_s J_{sk}^{p,q}$ for a streamer
 $(k_x p_{ti}, k_y p_{ti}) = (0, 4.4)$ at $t = 60-80 R/v_{ti}$




At the steady state,
normal cascade
dominates via the
direct coupling with
ion-scale turbulent
eddies.




After suppression of ETG/streamers, normal cascade dominates electron scales.

Triad transfer $\sum_s J_{sk}^{p,q}$ for a streamer
 $(k_x p_{ti}, k_y p_{ti}) = (0, 4.4)$ at $t = 60-80 R/v_{ti}$




At the steady state,
normal cascade
dominates via the
direct coupling with
ion-scale turbulent
eddies.




After suppression of ETG/streamers, normal cascade dominates electron scales.

Triad transfer $\sum_s J_{sk}^{p,q}$ for a streamer
 $(k_x p_{ti}, k_y p_{ti}) = (0, 4.4)$ at $t = 60-80 R/v_{ti}$



At the steady state,
normal cascade
dominates via the
direct coupling with
ion-scale turbulent
eddies.





Inefficient zonal mode generation in multi-scale turbulence.

- Inverse cascade from electron to ion scales seems not to be responsible.

Comparing multi-scale simulation with single-scale one,

- Zonal part of field energy is relatively weak.
- Inefficient zonal mode generation is observed.





Enhancement of ion-scale turbulence is caused by damping of zonal modes.

Splitting ion- and electron-scale contributions clarifies

- Electron-scale turbulence has damping effects on zonal modes around $k_x p_{ti} \sim 1$.
- The reduction of ZF shearing enhances transport.

Entropy transfer spectrum




Summary and discussion

We have first analyzed multi-scale ITG/TEM/ETG turbulence with **real mass ratio** and **β value**.

- We have demonstrated **the existence of multi-scale interactions even with real mass ratio.**
 - ETG/Streamers are suppressed by ITG/TEM turbulence.
- **ITG stabilization by finite- β effects** makes electron-scale contributions non-negligible.
 - We newly found that **electron-scale turbulence can enhance ion-scale turbulent transport.**

Summary and discussion

In terms of turbulence physics




- ✓ Normal cascade via coupling with ITG/TEM turbulent eddies dominates electron scales.
- ✓ Electron-scale turbulence acts as effective damping of ZFs.

In terms of transport level estimation

- Multi-scale spectrum is NOT a sum of single scales.
- When ITGs are highly unstable, ion-scale simulations give a good estimate of transport levels.
- In high- β regimes, electron scales can be important.
 - Effective damping of ion-scale ZFs


Multi-scale turbulence simulation ($\beta=2.0\%$)

Time evolution of the electrostatic potential fluctuations



Multi-scale turbulence simulation ($\beta=2.0\%$)

Time evolution of the electrostatic potential fluctuations



Multi-scale turbulence simulation ($\beta=2.0\%$)

Time evolution of the electrostatic potential fluctuations

