



Contribution ID: 342

Type: Oral

Role of MHD dynamo in the formation of 3D equilibria in fusion plasmas

Tuesday, 18 October 2016 10:45 (20 minutes)

This work investigates the formation of helical core equilibria in toroidal fusion plasmas, focusing on the role of dynamo, or magnetic flux pumping mechanisms in determining the equilibrium current profile. Dynamo effects determine the safety factor profile of the final 3D equilibrium, with important consequences on MHD stability and transport. We compare experimental results from multiple machines (RFX-mod, MST, AUG, DIII-D) and nonlinear MHD modelling. Two paradigmatic cases of helical state formation are considered and common physics is identified, by direct measurements of dynamo effects and MHD simulations: spontaneous formation in high-current reversed-field pinch (RFP) plasmas [1] and the hybrid scenario in high-beta tokamak plasmas [2]. Helical cores form in both cases, either spontaneously via saturation of MHD modes, or due to the marginally-stable ideal MHD response to external 3D fields. Direct measurements of the dynamo emf associated to 3D plasma distortions will be presented for a database of helical RFP plasmas from RFX-mod and MST, covering a wide range of plasma parameters. Similar measurements were also done in helical states forming in response to external 3D fields in Ohmic RFX-mod tokamak plasmas and in DIII-D high-beta hybrid plasmas. Experimental results qualitatively agree with nonlinear MHD modelling performed with the codes SpeCyl [3], PIXIE3D [4], and NIMROD [5]. They indicate that central current is redistributed by a dominantly electrostatic MHD dynamo. The underlying physics common to RFP and tokamak is thus revealed: a helical core displacement modulates parallel current density along flux tubes, which requires a helical electrostatic potential to build up, giving rise to a helical dynamo flow. Similar results were also recently obtained with the M3D-C1 code [6].

[1] R. Lorenzini et al., *Nature Phys.* 5, 570 (2009); [2] T.C. Luce et al., *Nucl. Fusion* 54, 013015 (2014); [3] D. Bonfiglio et al., *Phys. Rev. Lett.* 94, 145001 (2005); [4] D. Bonfiglio et al., *Plasma Phys. Control. Fusion* 57, 044001 (2015); [5] J.R. King, C.R. Sovinec, V.V. Mirnov, *Phys. Plasmas* 19, 055905 (2012); [6] S.C. Jardin et al., *Phys. Rev. Lett.* 115, 215001 (2015).

Paper Number

EX/1-1

Country or International Organization

Italy

Primary author: PIOVESAN, Paolo (Consorzio RFX)

Co-authors: Dr ZANIOL, Barbara (Consorzio RFX, Padova, Italy); Dr PAZ-SOLDAN, Carlos (General Atomics, San Diego, CA, USA); Dr HOLCOMB, Christopher T. (Lawrence Livermore National Laboratory); Dr CHRYSTAL, Colin (Oak Ridge Associated Universities, Oak Ridge, TN, USA); Dr BONFIGLIO, Daniele (Consorzio RFX, Padova, Italy); Dr TERRANOVA, David (Consorzio RFX, Padova, Italy); Dr ESCANDE, Dominique (Laboratoire PIMM, UMR

7345 CNRS-Aix Marseille Université, France); Dr TURCO, Francesca (Columbia University); Prof. SARFF, John (University of Wisconsin-Madison); Dr PIRON, Lidia (CCFE, Abingdon, UK); Dr MARRELLI, Lionello (Consorzio RFX); Dr CHACÓN, Luis (Los Alamos National Laboratory, Los Alamos, NM, USA); NORNBERG, Mark (University of Wisconsin-Madison); Dr TAYLOR, N.Z. (Oak Ridge Associated Universities, Oak Ridge, TN, USA); Dr FRANZ, Paolo (Consorzio RFX, Padova, Italy); Dr ZANCA, Paolo (Consorzio RFX, Padova, Italy); Dr CAPPELLO, Susanna (Consorzio RFX); Dr LUCE, Timothy C. (General Atomics); Dr IGOCHINE, Valentin (MPI für Plasma-physik, Garching, Germany)

Presenter: PIOVESAN, Paolo (Consorzio RFX)

Session Classification: 3D Physics

Track Classification: EXS - Magnetic Confinement Experiments: Stability