



IAEA FEC 201

Contribution ID: 291

Type: Oral

Penetration and amplification of resonant perturbations in 3D ideal-MHD equilibria

Tuesday, 18 October 2016 11:25 (20 minutes)

The nature of ideal-MHD equilibria in three-dimensional geometry is profoundly affected by resonant surfaces, which beget a non-analytic dependence of the equilibrium on the boundary. Furthermore, non-physical currents arise in equilibria with continuously-nested magnetic surfaces and smooth pressure and rotational-transform profiles.

We demonstrate that three-dimensional, ideal-MHD equilibria with nested surfaces and delta-function current-densities that produce a discontinuous rotational-transform are well defined and can be computed both perturbatively and using fully-nonlinear equilibrium calculations.

The results are of direct practical importance: we predict that resonant magnetic perturbations penetrate past the rational surface (i.e. “shielding” is incomplete, even in purely ideal-MHD) and that the perturbation is amplified by plasma pressure, increasingly so as stability limits are approached.

Paper Number

TH/1-1

Country or International Organization

USA

Primary author: Dr HUDSON, Stuart (Princeton Plasma Physics Laboratory)

Co-author: Dr LOIZU, Joaquim (Max Planck Institut for Plasma Physic)

Presenter: Dr HUDSON, Stuart (Princeton Plasma Physics Laboratory)

Session Classification: 3D Physics

Track Classification: THS - Magnetic Confinement Theory and Modelling: Stability