



IAEA FEC 2011

Contribution ID: 90

Type: Poster

Fast heating of an imploded core under counter beam irradiation by using a repetitive IFE driver HAMA

Thursday, 20 October 2016 08:30 (4 hours)

Fast heating is a scheme to heat an imploded high density plasma into a hot state before the plasma hydrodynamically expands and dissolves. For a heating process driven by laser induced hot electron currents, the energy of hot electrons beyond MeV having a range of a few g/cm^2 , therefore collisional coupling is inefficient until the core density reaches hundreds g/cc . To improve the MeV electrons coupling, an energy deposition via the anomalous resistivity in strong magnetic filaments induced by the electromagnetic tow-stream instability, so called Weibel instability was expected. However, this Weibel instability was found to be suppressed in the collisional dense plasma, namely, the heating by the fast electrons stays inefficient. In contrast, counter irradiating fast heating scheme we propose here can improve this insufficient coupling efficiency by introducing a magnetic field supposed to be contributing to trap hot electrons in the core. We report fast heating of a shock-imploded core under counter beam irradiation by using a repetitive IFE driver HAMA. Experimental results show that (i) a shock-imploded core with 70 micron diameter, originally deuterated polystyrene (CD) spherical shell of 500 micron diameter, is flashed by counter irradiating 110 fs, 7 TW laser pulses. The coupling efficiency from the laser to the core emission was inferred 13%. A collisional particle-in-cell (PIC) simulation indicates a possibility that counter hot electron currents generate magnetic filaments in the imploded core. (ii) Fast electrons with energy bellow a few MeV might be trapped by these filaments in the core region supposed to be contributing to the observed X-ray flash and the high coupling efficiency. These results indicate a possibility that counter irradiating fast heating scheme can improve the energy coupling into the core by hot electrons those are limited to less 10% for one-side irradiation fast heating conducted so far.

Paper Number

IFE/P5-9

Country or International Organization

Japan

Primary author: Prof. MORI, Yoshitaka (The Graduate School for the Creation of New Photonics Industries)

Co-authors: Dr SUNAHARA, Atsushi (Institute for Laser Technology); Dr MIURA, Eisuke (AIST); Dr KAN, Hirofumi (Hamamatsu Photonics, K. K.); Dr AZUMA, Hirozumi (Aichi SR); Prof. ISHII, Katsuhiko (GPI); Mr SATO, Nakahiro (Hamamatsu Photonics, K. K.); Dr KOMEDA, Osamu (Toyota Technical Development Corp.); Prof. HANAYAMA, Ryohei (The Graduate School for the Creation of New Photonics Industries); Mr NAKAYAMA, Suisei (GPI); Dr KURITA, Takashi (Hamamatsu Photonics, K. K.); Mr SEKINE, Takashi (Hamamatsu Photonics)

K.K.); Dr HIOKI, Tatsumi (GREMO, Nagoyga Univ.); Mr NISHI, Teppei (TOYOTA Central Research and Development Laboratories, Inc.); Prof. MOTOHIRO, Tomoyoshi (GREMO, Nagoyga Univ.); Dr KAWASHIMA, Toshiyuki (Hamamatsu Photonics, K. K.); Mr NISHIMURA, Yasuhiko (The Graduate School for the Creation of New Photonics Industries, Toyota Technical Development Corporation); Prof. SENTOKU, Yasuhiko (Univ of Nevada, Reno); Prof. KITAGAWA, Yoneyoshi (The Graduate School for the Creation of New Photonics Industries)

Presenter: Prof. MORI, Yoshitaka (The Graduate School for the Creation of New Photonics Industries)

Session Classification: Poster 5

Track Classification: IFE - Inertial Fusion Experiments and Theory