Contribution ID: **945** Type: **Poster**

The effect of the isotope on the H-mode density limit

Friday, 21 October 2016 14:00 (4h 45m)

Identification of the mechanisms for the H-mode density limit in machines with fully metallic walls, and their scaling to future devices is essential to find for these machines the optimal operational boundaries with the highest attainable density and confinement. Systematic investigations of H-mode density limit plasmas in experiments with deuterium and hydrogen external gas fuelling have been performed on JET machine with fully metallic walls, and results have been compared with one another.
The observed H-mode density limit on JET in D- as well as in H-plasmas demonstrates similar operation phases: the stable H-mode phase, degrading H-mode, breakdown of the H-mode with energy confinement deterioration accompanied by a dithering cycling phase, followed by the L-mode phase. Independent of the isotopic effect, total radiated power as well as the radiation power in the main chamber (Prad, bulk) stays almost constant during the H-mode phase until the H-L transition. The density limit is not related to an inward collapse of the hot discharge core induced by overcooling of the plasma periphery by radiation. It was observed in D- and H-plasmas that detachment, as well as the X-point MARFE itself, does not trigger the H-L transition and thus does not present a limit on the plasma density and that it is the plasma confinement, most likely determined by edge parameters, which is ultimately responsible for the H-mode DL. Independent of the isotopic mas of the main plasma, it has been observed that the transition from H-mode to L-mode is not always an abrupt event but may exhibit a series of H-L-H transitions ('dithering H-mode"), or a gradual transition (which is orders of magnitude longer than energy confinement time τE). Although the operation phases are identical for D- and H-plasmas, the DL shows strong dependency on the isotopic mass effect, the DL is up to 40% lower in the H-plasma than in the deuterium plasma. Basically, the density limit in H mode on JET-ILW is nearly independent of the power in the range of observed heating powers and the corresponding densities only approach in this configuration a Greenwald fraction of about fGW=0.9 and fGW=0.84 in D-plasma and in H-plasma correspondingly.
The measured Greenwald fractions are found to be consistent with the predictions from theoretical model based on MHD instability theory in the near-SOL.

Paper Number

PDP-8

Country or International Organization

Germany

Primary author: Dr HUBER, Alexander (Forschungszentrum Jülich GmbH, Institut für Energie- und Klimaforschung –Plasmaphysik, Partner of the Trilateral Euregio Cluster (TEC), 52425 Jülich, Germany)

Co-authors: Dr SIPS, Adrianus (EFDA-JET); Dr CHANKIN, Alex (Max-Planck-Institut für Plasmaphysik, D-85748 Garching, Germany); Dr BOBOC, Alexandru (CCFE, Culham Science Centre, Abingdon, OX14 3DB, UK); Dr MEIGS, Andrew (CCFE, Culham Science Centre, Abingdon, OX14 3DB, UK); Dr KALLENBACH, Arne

(Max-Planck-Institut f. Plasmaphysik); Prof. LINSMEIER, Christian (Forschungszentrum Jülich GmbH, Institut für Energie- und Klimaforschung –Plasmaphysik, Partner of the Trilateral Euregio Cluster (TEC), 52425 Jülich, Germany); Dr GUILLEMAUT, Christophe (Instituto de Plasmas e Fusão Nuclear, Instituto Superior Técnico, Universidade Lisboa, PT); Dr LOWRY, Christopher (European Commission); Dr MAGGI, Costanza (CCFE); Dr CARRALERO, Daniel (IPP Garching); Dr VIEZZER, Eleonora (Max-Planck-Institut fuer Plasmaphysik); Dr JOFFRIN, Emmanuel (CEA/IRFM); Dr DELABIE, Ephrem (EURATOM-FOM association DIFFER , The Netherlands); Dr REIMOLD, Felix (Forschungszentrum Jülich); Dr SERGIENKO, Gennady (Forschungszentrum Jülich GmbH,); Mr MATTHEWS, Guy (Culham Centre for Fusion Energy); Dr ESSER, Hans Guenter (Forschungszentrum Jülich GmbH, Institut für Energie- und Klimaforschung –Plasmaphysik, Partner of the Trilateral Euregio Cluster (TEC), 52425 Jülich, Germany); Prof. ZOHM, Hartmut (Max-Planck-Institut für Plasmaphysik); Dr SCHWEINZER, Josef (IPP Garching); Dr WISCHMEIER, Marco (IPP Garching); Dr BRIX, Mathias (CCFE, Culham Science Centre, Abingdon, OX14 3DB, UK); Dr BERNERT, Matthias (Max-Planck-Institut für Plasmaphysik, D-85748 Garching, Germany); Dr STAMP, Mike (CCFE, Culham Science Centre, Abingdon, OX14 3DB, UK); Mr ABREU, Paulo (Instituto de Plasmas e Fusão Nuclear, Instituto Superior Técnico, Universidade Lisboa, PT); Dr LANG, Peter (Max-Planck-Institut fűr Plasmaphysik); Dr MERTENS, Philippe (Forschungszentrum Juelich, EURATOM-Association); Dr BREZINSEK, Sebastijan (Forschungszentrum Jülich); Mr JACHMICH, Stefan (Laboratory for Plasma Physics, ERM/KMS, B-1000 Brussels, Belgium); Dr WIESEN, Sven (Forschungszentrum Jülich GmbH, Institut für Energie- und Klimaforschung –Plasmaphysik, Partner of the Trilateral Euregio Cluster (TEC), 52425 Jülich, Germany); Dr EICH, Thomas (Max-Planck-Institute for Plasma Physics); Dr KRUEZI, Uron (5CCFE, Culham Science Centre, Abingdon, OX14 3DB, UK); Mrs HUBER, Valentina (Forschungszentrum Jülich GmbH, Supercomputing Centre, 52425 Jülich, Germany)

Presenter: Dr WIESEN, Sven (Forschungszentrum Jülich GmbH, Institut für Energie- und Klimaforschung – Plasmaphysik, Partner of the Trilateral Euregio Cluster (TEC), 52425 Jülich, Germany)

Session Classification: Poster 8

Track Classification: EXD - Magnetic Confinement Experiments: Plasma–material interactions; divertors; limiters; scrape-off layer (SOL)