



Contribution ID: 133

Type: **Poster**

## Excitation of frequency jump by barely Passing Electrons

*Tuesday, 18 October 2016 08:30 (4 hours)*

An e-fishbone frequency jump has been observed on Tore Supra, which is important for the redistribution of energetic electrons and energetic particle losses. E-fishbone periodic frequency jump phenomena are also observed on HL-2A. Soft X-ray tomography shows that the poloidal and toroidal mode numbers are 1/1 and 2/2 with the frequency jump. In this paper we present a theoretical base of the frequency jump in the e-fishbone experiments. It is identified that barely passing electrons are the drive of the e-fishbone, rather than the trapped electrons. The frequency jumps in HL-2A E-fishbone experiments are numerically reproduced. E-fishbone frequency increases with the hot electron energy which is consistent with the experiments. The growth rate of the mode ( $m=2, n=2$ ) is greater than the one of the mode ( $m=1, n=1$ ) in contrast to the pure MHD prediction. The calculated temporal evolutions of the hot electron energy and the kink mode amplitude are periodic which in good agreement with the observed e-fishbone jump cycle. The theory provides an insight on HL-2A and Tore Supra experiments.

### Paper Number

TH/P1-32

### Country or International Organization

China

**Primary author:** Prof. FENG, Hao (School of Science, Xihua University)

**Co-authors:** Dr LI, Huidong (School of Science, Xihua university); Prof. WANG, Zhongtian (School of Science, Xihua University;Southwestern Institute of Physics)

**Presenter:** Prof. WANG, Zhongtian (School of Science, Xihua University;Southwestern Institute of Physics)

**Session Classification:** Poster 1

**Track Classification:** THS - Magnetic Confinement Theory and Modelling: Stability