



Contribution ID: 409

Type: Poster

Gyrokinetic simulations of an electron temperature gradient turbulence-driven current in tokamak plasmas

Tuesday, 18 October 2016 14:00 (4h 45m)

The so-called “spontaneous” or “intrinsic” rotation driven by ion-scale turbulence has been widely observed in tokamaks. If we turn our attention to the electron parallel momentum balance, it is likely that electron-scale turbulence, e.g. electron temperature gradient (ETG) turbulence, can modify the Ohm’s law, hence providing a current source. However, there has been no serious study of an ETG-driven current in self-consistent simulations using realistic tokamak geometry. In this work, we report results of a gyrokinetic simulation study elucidating the characteristics of an intrinsic current driven by ETG turbulence in toroidal geometry. We focus on effects of the normalized electron gyroradius ρ_{e0} on the ETG-driven current. Our simulations demonstrate that the amount of the ETG-driven current increases with ρ_{e0} , as expected from the gyro-Bohm scaling. In particular, a perturbation of a q-profile by the ETG-driven current becomes visible when $a < 4000 \rho_{e0}$. This finding suggests that a significant intrinsic current can be driven inside an H-mode pedestal where the steep gradient of an electron temperature pedestal can excite ETG turbulence in a narrow region.

Paper Number

TH/P2-5

Country or International Organization

Republic of Korea

Primary author: Dr YI, Sumin (National Fusion Research Institute, Daejeon, Republic of Korea)

Co-authors: Dr JHANG, Hogun (National Fusion Research Institute, Daejeon, Republic of Korea); Dr KWON, Jae-Min (National Fusion Research Institute, Daejeon, Republic of Korea)

Presenter: Dr YI, Sumin (National Fusion Research Institute, Daejeon, Republic of Korea)

Session Classification: Poster 2

Track Classification: THC - Magnetic Confinement Theory and Modelling: Confinement