



IAEA FEC 201

Contribution ID: 659

Type: Poster

Long-lived pressure-driven MHD mode in KSTAR plasmas

Wednesday, 19 October 2016 14:00 (4h 45m)

Highly coherent structures associated with an extremely long-lived saturated magnetohydrodynamic (MHD) instability have been observed in KSTAR tokamak under a long-pulse and steady-state operation. As the plasma performance is increased in advanced tokamak regimes, possible deleterious effects of MHD modes become more important, especially for steady-state burning plasmas in the next-step devices such as ITER and DEMO. One of the commonly seen modes is the $m = 2$, $n = 1$ resistive kink mode that either leads to confinement degradation, or mode locking followed by a full disruption. In KSTAR, however, long-pulse discharges regularly exhibit a coherent structure in the form of a saturated pressure-driven MHD mode that can be sustained as long as 40 seconds, the full discharge duration, when the mode is located near the plasma core region with a broad safety factor profile with q_0 larger than 2.

Paper Number

EX/P4-10

Country or International Organization

Korea

Primary author: Dr LEE, Sang Gon (National Fusion Research Institute)

Co-authors: Dr AYDEMIR, Ahmet (National Fusion Research Institute, Daejeon, Korea); Dr YOUNG SOON, Bae (National Fusion Research Institute); Mr CHOE, G. H. (POSTECH); Prof. YUN, Gunsu (Pohang University of Science and Technology); Dr LEE, Hyungho (National Fusion Research Institute); Dr SEOL, JaeChun (National Fusion Research Institute); Ms YOO, Jeong Won (National Fusion Research Institute); Dr BAK, Jun Gyo (National Fusion Research Institute); Dr TERZOLO, Laurent (National Fusion Research Institute); Mr KYU DONG, Lee (National Fusion Research Institute)

Presenter: Dr LEE, Sang Gon (National Fusion Research Institute)

Session Classification: Poster 4

Track Classification: EXW - Magnetic Confinement Experiments: Wave-plasma interactions; current drive; heating; energetic particles