



IAEA FEC 201

Contribution ID: 123

Type: Poster

## Effect of the Transition to Improved Core Confinement Observed in the LHCD Experiment at FT-2 Tokamak

Friday, 21 October 2016 08:30 (4 hours)

To explain a relatively good efficiency of LHCD and improved core confinement transition obtained at the small FT-2 tokamak ( $R = 0.55m$ ,  $a = 0.08m$ ,  $B_T \leq 3T$ ,  $I_{pl} = 35kA$ ,  $f_0 = 920MHz$ ,  $\Delta t_{pl} = 50ms$ ,  $\Delta t_{RF}$  = from 30ms to 36ms) [1] a thorough modeling of experimental data has been performed. Effect of LHW on the transition into improved core confinement regime is discussed in the deuterium plasma experiment. It was observed, that in the LHCD experiment with initial OH density  $\langle n_e \rangle = 1.6 \times 10^{19} m^{-3}$  the central electron temperature  $T_e(r = 0 \text{ cm})$  measured by TS diagnostics increases during RF pulse from 550eV to 700eV and that is accompanied by cooling of the plasma periphery and the density rise. This effect could not be explained by increase of working gas or impurity recycling because the  $D\beta$  line intensity and radiation losses during RF pulse is not appreciably changed.

According to GRILL3D, FRTC and ASTRA codes modeling the increase of the density and electron temperature  $T_e$  inside of  $r < 3cm$  (despite the decrease of ohmic heating power POH at LHCD) happens due to strong reduction of the electron transport in this region where the magnetic shear vanishes, and the value of thermal diffusivity  $\chi_{e,eff}$  decreases. Broadening of the plasma current profile by noninductive LHCD results in flattening of the safety factor  $q$  - profile in the plasma column center. As the result, the magnetic shear  $s = (r/q)(dq/dr)$  in the center became low, or even negative. In such a case the transport code (where the electron transport was described by the mixed Bohm and gyro-Bohm model) predicts a reduction of the transport [2]. Paper presents new experimental data and modeling results appropriate to the transition to improved core confinement during LHCD experiment. In particular, special attention one attends to the experimental periphery data and data of the threshold power for transition to improved core confinement in deuterium/hydrogen plasma.

This work (for S.I. Lashkul and FT-2 team) was supported in part by the Russian Foundation for Basic Research project nos 14-08-00476

References

- [1] S.I. Lashkul, A.B. Altukhov, A.D. Gurchenko, E.Z. et al. Nucl. Fusion 55 (2015) 073019
- [2] Y. Peysson and the TORE SUPRA Team. 2000 Plasma Phys. Control. Fusion 42 B87–B114

### Paper Number

EX/P7-41

### Country or International Organization

Russian Federation

**Primary author:** Dr LASHKUL, Sergey (Ioffe Institute)

**Co-authors:** Dr SAVELIEV, Alexandr (Ioffe Institute); Dr ALTUKHOV, Alexey (Ioffe Institute); Dr GURCHENKO, Alexey (Ioffe Institute); Mr PEREVALOV, Arture (St. Petersburg State Polytechnical University); Dr KOUPRIENKO, Denis (Ioffe Institute); Prof. GUSAKOV, Evgeniy (Ioffe Institute); Dr IRZAK, Michail (Ioffe Institute); Dr KANTOR, Michail (Ioffe Institute); Dr SHATALIN, Sergey (St. Petersburg State Polytechnical University); Dr DYACHENKO, Valeriy (Ioffe Institute)

**Presenter:** Dr LASHKUL, Sergey (Ioffe Institute)

**Session Classification:** Poster 7

**Track Classification:** EXC - Magnetic Confinement Experiments: Confinement