



IAEA FEC 2016

Contribution ID: 578

Type: Poster

Nonlinear dynamics of ELMs with Er shear and collisionality trends

Thursday, 20 October 2016 14:00 (4h 45m)

Simulations with the BOUT++ code have been used to study the energy loss for edge localized modes (ELMs) at different collisionality & electric field E_r shear and to investigate the controls of transition to different ELM and quiescent H-mode (QH-mode) regimes. The simulation results indicate that by development of a flexible E_r control capabilities, it is conceivable that tokamak operation regime access can be achieved through a control of edge fluctuation spectrum via the radial electric field and its shear. By decreasing collisionality with a increasing E_r for a narrow pedestal, nonlinear simulations show that (1) power spectrum becomes narrower and linear growth rate increases, the dominant mode decreasing from high- n ballooning modes to low- n peeling modes; (2) Bispectrum analysis shows that nonlinear mode coupling becomes weaker, resulting in the dominant filamentary structures and increasing ELM energy loss. The increasing E_r shear at high collisionality with a narrow pedestal leads to strong nonlinear coupling and reduced ELM energy loss. In contrast, the increasing E_r shear at low collisionality with a wide pedestal can modify the stability boundary to drive stable low- n peeling modes unstable, leading to enhanced pedestal transport due to either a saturated low- n peeling modes or a broadband turbulence, therefore leading to a reduced pedestal pressure gradient, allowing the development of a broader and thus higher transport barrier in QH mode without ELMs.

To validate BOUT++ simulations against experimental data for inter-ELM fluctuations, the BOUT++ simulations are performed based on a set of C-Mod and DIII-D experiment data, the overall signatures of simulation results for quasi-coherent fluctuations (QCF) show good agreement with C-Mod and DIII-D measurements. QCFs are localized in the pedestal region having a predominant frequency at $f \approx 300\text{--}400\text{kHz}$ and poloidal wavenumber at $k_{\theta} \approx 0.7/\text{cm}$, and propagate in the electron diamagnetic direction in the laboratory frame. (2) The pedestal profiles giving rise to QCFs are near the marginal instability threshold for ideal peeling-ballooning (P-B) modes for both C-Mod and DIII-D. (3) Particle diffusivity is either smaller than the heat diffusivity for DIII-D or similar to the heat diffusivity for C-Mod.

This work was performed for USDOE by LLNL under DE-AC52-07NA27344.LLNL-ABS-680566.

Paper Number

TH/P6-18

Country or International Organization

USA

Primary author: XU, Xueqiao (Lawrence Livermore National Laboratory)

Co-authors: Dr DIALLO, Ahmed (PPPL); Mr MA, Chenhao (Peking University); Dr KONG, Defeng (Institute of Plasma Physics Chinese Academy of Sciences); HUGHES, Jerry (MIT PSFC); Mr CHEN, Jianguo (Peking University); Dr YAN, Ning (Institute of Plasma Physics, Chinese Academy of Sciences, Hefei, China); Prof. DIA-

MOND, Patrick (University of California San Diego); Dr SNYDER, Philip B. (General Atomics); Mr GROEBNER, R.J. (General Atomics)

Presenter: XU, Xueqiao (Lawrence Livermore National Laboratory)

Session Classification: Poster 6

Track Classification: THD - Magnetic Confinement Theory and Modelling; Plasma-material interactions; divertors, limiters, SOL