



IAEA FEC 2016

Contribution ID: 736

Type: Poster

## ELM Characterization and Dynamics at Near-Unity $A$ in the Pegasus ST

Wednesday, 19 October 2016 14:00 (4h 45m)

Operation in the high confinement (H-mode) regime and mitigation of associated deleterious Edge Localized Mode (ELM) activity are necessary for the success of ITER and future reactors. H-mode studies at near-unity aspect ratio  $A$  can offer unique insights into these issues. Edge plasma parameters at low  $A$  permit unique measurements of the edge pedestal with high spatiotemporal resolution using probes. In particular, measurements of the current density profile  $J_{\text{edge}}$  of import to peeling-ballooning stability and its nonlinear dynamics during ELMs are presented. Two classes of ELMs have been identified to date by their proximity to  $P_{\text{LH}}$  and measured  $n$  spectra provided by a near-edge Mirnov coil array. Both ELM types produce propagating, field-aligned filaments and have multiple  $n$  measured during the crash. These observations are consistent with the presence of a spectrum of simultaneously unstable peeling-ballooning modes anticipated by theory and nonlinear ELM simulations. Small, Type III-like ELMs occur at  $P_{\text{OH}} \sim P_{\text{LH}}$  with  $n \leq 4$ . Large, Type-I-like ELMs occur with  $P_{\text{OH}} > P_{\text{LH}}$  and intermediate  $5 < n < 15$ , similar to ELMs at  $A \sim 1.3$  in NSTX. The Type III  $n$  ranges are opposite that reported at high  $A$ , and Type I  $n$  are in the low range of those reported at high  $A$ . These differences presumably reflect the strong peeling mode drive  $\sim J_{\text{edge}}/B$  present in the ST. The dominant  $n$  component of a large ELM grows exponentially, whereas other  $n$  are nonlinearly driven and damped prior to the crash. Access to small and large ELMs are demonstrated in Ohmic H-mode plasmas by varying the applied input power.  $J_{\text{edge}}(R,t)$  measurements have been obtained across single ELM events with sub-cm spatial and Alfvénic temporal resolution. Both ELM types feature the nonlinear generation of “current-hole”  $J_{\text{edge}}$  perturbations, similar to prior studies of nonlinear peeling mode dynamics in Pegasus. A Type I ELM is shown to additionally expel a current-carrying filament during the ELM crash. Initial experiments coupling small amounts of helical edge current injection to H-mode plasmas suggest suppression of Type III ELM activity and negligible macroscopic impact on the discharge. This occurs for injected currents  $I_{\text{inj}} < 1$  kA. Above  $I_{\text{inj}} > 1$  kA the 3D field perturbation degrades the edge sufficiently to exit H-mode.

### Paper Number

EX/P4-51

### Country or International Organization

United States of America

**Primary author:** Dr BONGARD, Michael (University of Wisconsin-Madison)

**Co-authors:** Mr KRIETE, David (University of Wisconsin-Madison); Mr BARR, Jayson (University of Wisconsin-Madison); Dr REUSCH, Joshua (University of Wisconsin-Madison); Dr THOME, Kathreen (University of Wisconsin-Madison); Prof. FONCK, Raymond (University of Wisconsin-Madison)

**Presenter:** Dr BONGARD, Michael (University of Wisconsin-Madison)

**Session Classification:** Poster 4

**Track Classification:** EXW - Magnetic Confinement Experiments: Wave-plasma interactions; current drive; heating; energetic particles