



Contribution ID: 535

Type: **Poster**

Equilibrium solutions of MHD equations for GAMs in the edge tokamak plasma

Tuesday, 18 October 2016 08:30 (4 hours)

Numerical calculations of nonlinear MHD equations in frames of reduced two-fluid Braginskij equations for geodesic acoustic modes (GAM) with $n = 0$, $m = 0, +1, -1$ in high collisional edge tokamak plasma were performed. $N = 0$, $m = 0, +1, -1$. It was shown that with account of parallel dissipation (finite conductivity σ_{\parallel}) allows us to obtain the steady state equilibrium solutions for GAMs. The obtained 2D equilibrium includes the velocity of poloidal rotation and the equilibrium electric potential, which value is close to well-known Pfirsch-Schluter potential. It was shown that the main role in formation of the equilibrium poloidal rotation plays two forces: the Stringer-Winsor force and the neoclassical force, linked with the parallel viscosity. Maximum values of GAM are located near the maximum of pressure gradient. Calculated radial profile of electric field E looks like the parabolic negative well ($E < 0$).

Paper Number

TH/P1-31

Country or International Organization

Russian Federation

Primary author: Dr SHURYGIN, Remir (NRC 'Kurchatov Institute', Moscow, Russia)

Co-author: Dr MELNIKOV, Alexander (NRC 'Kurchatov Institute')

Presenter: Dr SHURYGIN, Remir (NRC 'Kurchatov Institute', Moscow, Russia)

Session Classification: Poster 1

Track Classification: THS - Magnetic Confinement Theory and Modelling: Stability