



IAEA FEC 2016

Contribution ID: 842

Type: Poster

Mitigation of Runaway Current with Supersonic Molecular Beam Injection on HL-2A Tokamak

Friday, 21 October 2016 08:30 (4 hours)

Disruption mitigation experiments with MGI have been carried out on HL-2A with a rapid (~ 1 ms), massive ($\sim 10^{21}$ particles) injection of helium/neon/argon to study various injection scenarios. The behaviors of runaway currents in MGI induced disruptions have been investigated. A long-lasting RE plateau is achieved after argon injection by MGI even at $B_t = 1.28$ T, much lower than previous B_t threshold found in other tokamaks. It was found that argon injection can cause the generation of runaways carrying up to 30% of the initial plasma current, while disruptions triggered by injection of helium or neon are runaway free. Furthermore, the runaway current caused by argon injection with MGI was successfully suppressed by SMBI with a number of injected atoms of about 1.0×10^{21} . Light gases, such as helium, are selected in this experiment for its high efficiency in increasing the density. These experiments suggest that SMBI might be viable for runaway suppression in future tokamaks even though core penetration of jet neutrals is not achieved. An understanding of this paradox is obtained by modeling, which shows that the initial cooling of the plasma periphery triggers a very rapid growth of low-order tearing mode, resulting in a stochastic region over much of the plasma. This allows rapid transport across the entire plasma, and could explain the effectiveness of SMBI mitigation in HL-2A in spite of the shallow penetration of the neutral gas jet. In addition, a toroidal alfvén eigenmode (TAE) was observed during disruptions deliberately triggered by the massive gas injection (MGI) of argon. This mode occurs at the beginning of the current quench and lasts about 1-2 ms. These instabilities appear to be favorable in limiting the RE beam formation. It has been found that the runaway plateau is easy to obtain on the condition of high normalized magnetic fluctuation level ($\delta B/B$), the runaway plateau is even invisible when $\delta B/B$ exceeds the threshold of about 7.8×10^{-4} , indicating that this magnetic mode plays a scattering role on the RE beam strength.

Paper Number

EX/9-3

Country or International Organization

China

Primary author: Prof. LIU, Yi (southwestern institute of physics)

Presenter: Prof. LIU, Yi (southwestern institute of physics)

Session Classification: Poster EX/9, EX/10, TH/7, TH/8, IFE/1, MPT/1, FNS/1

Track Classification: EXS - Magnetic Confinement Experiments: Stability